

Imfinzi® (durvalumab) (Intravenous)

Document Number: IC-0301

Last Review Date: 03/02/2023

Date of Origin: 05/30/2017

Dates Reviewed: 05/2017, 08/2017, 11/2017, 02/2018, 05/2018, 09/2018, 12/2018, 03/2019, 06/2019, 09/2019, 12/2019, 03/2020, 06/2020, 09/2020, 12/2020, 03/2021, 04/2021, 06/2021, 09/2021, 12/2021, 03/2022, 06/2022, 09/2022, 10/2022, 12/2022, 03/2023

I. Length of Authorization ^{Δ 1}

Coverage will be provided for 6 months and may be renewed (unless otherwise specified).

- Non-Small Cell Lung Cancer (single agent use as consolidation therapy): Coverage will be provided for 6 months and may be renewed up to a maximum of 12 months of therapy.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Imfinzi 120 mg/2.4 mL single-dose vial: 4 vials per 14 days
- Imfinzi 500 mg/10 mL single-dose vial: 2 vials per 14 days

B. Max Units (per dose and over time) [HCPCS Unit]:

- NSCLC: 112 billable units (1,120 mg) every 14 days
- SCLC: 150 billable units (1,500 mg) every 21 days x 6 doses, then 150 billable units (1,500 mg) every 28 days
- Biliary Tract Cancer: 150 billable units (1,500 mg) every 21 days x 8 doses, then 150 billable units (1,500 mg) every 28 days
- Hepatocellular Carcinoma: 150 billable units (1,500 mg) every 28 days
- Cervical Cancer: 150 billable units (1,500 mg) every 21 days x 4 doses, then 150 billable units (1,500 mg) every 28 days

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

Universal Criteria

- Patient has not received previous therapy with a programmed death (PD-1/PD-L1)-directed therapy (e.g., nivolumab, pembrolizumab, atezolizumab, avelumab, cemiplimab, dostarlimab, nivolumab/relatlimab-rmbw, etc.) unless otherwise specified^Δ; **AND**

Non-Small Cell Lung Cancer (NSCLC) † ‡ ^{1,3-5,16}

- Patient has unresectable stage II-III disease; **AND**
 - Patient has a performance status (PS) of 0-1; **AND**
 - Used as a single agent as consolidation therapy; **AND**
 - Disease has not progressed after definitive chemoradiation; **OR**
- Patient has recurrent, advanced, or metastatic disease; **AND**
 - Used as first-line therapy; **AND**
 - Used in one of the following:
 - Patients with tumors that are negative for actionable molecular biomarkers* and PD-L1 $\geq 1\%$ to 49%
 - Patients with PS of 0-1 who have tumors that are negative for actionable molecular biomarkers* and PD-L1 $< 1\%$
 - Patients with PS of 0-1 who are positive for one of the following molecular mutations: EGFR exon 20, KRAS G12C, BRAF V600E, NTRK1/2/3 gene fusion, MET exon-14 skipping, RET rearrangement, or ERBB2 (HER2); **AND**
 - Used in combination with tremelimumab-actl, albumin-bound paclitaxel, and carboplatin; **OR**
 - Used in combination with tremelimumab-actl, pemetrexed, and either carboplatin or cisplatin for nonsquamous cell histology; **OR**
 - Used in combination with tremelimumab-actl, gemcitabine, and either carboplatin or cisplatin for squamous cell histology; **OR**
 - Used as subsequent therapy; **AND**
 - Used in one of the following:
 - Patients with PS of 0-1 who are positive for one of the following molecular mutations: BRAF V600E, NTRK1/2/3 gene fusion, MET exon-14 skipping, or RET rearrangement
 - Patients with PS of 0-1 who are positive for one of the following molecular mutations and received prior targeted therapy§: EGFR exon 19 deletion or exon 21 L858R tumors, EGFR S768I, L861Q, and/or G719X mutation, ALK rearrangement, or ROS1 rearrangement; **AND**
 - Used in combination with tremelimumab-actl, albumin-bound paclitaxel, and carboplatin; **OR**
 - Used in combination with tremelimumab-actl, pemetrexed, and either carboplatin or cisplatin for nonsquamous cell histology; **OR**

- Used in combination with tremelimumab-actl, gemcitabine, and either carboplatin or cisplatin for squamous cell histology; **OR**
- Used as continuation maintenance therapy in patients who have achieved a tumor response or stable disease following initial therapy; **AND**
 - Used as a single agent following a first-line regimen with durvalumab and tremelimumab-actl plus chemotherapy; **OR**
 - Used in combination with pemetrexed following a first-line regimen with durvalumab, tremelimumab-actl, pemetrexed and either carboplatin or cisplatin for nonsquamous cell histology

** Note: Actionable molecular genomic biomarkers include EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET exon 14 skipping mutation, RET rearrangement, and ERBB2 (HER2). If there is insufficient issue to allow testing for all of EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET, RET, and ERBB2 (HER2) repeat biopsy and/or plasma testing should be done. If these are not feasible, treatment is guided by available results and, if unknown, these patients are treated as though they do not have driver oncogenes.*

Small Cell Lung Cancer (SCLC) † ‡ Ⓢ 1,3,7,8,10

- Patient has extensive stage disease (ES-SCLC); **AND**
 - Used as first-line therapy in combination with etoposide and either carboplatin or cisplatin; **OR**
 - Used as single-agent maintenance therapy after initial therapy with etoposide and either carboplatin or cisplatin

Hepatobiliary Cancers † ‡ Ⓢ 1,3,11,12,14,15

- Patient has hepatocellular carcinoma (HCC); **AND**
 - Used as first-line therapy as a single agent or in combination with tremelimumab-actl; **AND**
 - Patient has unresectable disease and is not a transplant candidate; **OR**
 - Patient has liver-confined disease that is inoperable by performance status, comorbidity, or with minimal or uncertain extrahepatic disease; **OR**
 - Patient has metastatic disease or extensive liver tumor burden; **OR**
- Patient has biliary tract cancer (e.g., gallbladder cancer or intra-/extra-hepatic cholangiocarcinoma); **AND**
 - Used in combination with cisplatin and gemcitabine; **AND**
 - Used as primary treatment for unresectable, locally advanced, or metastatic disease; **OR**
 - Used for recurrent disease >6 months after surgery with curative intent and >6 months after completion of adjuvant therapy

Cervical Cancer † ‡ 3,17

- Patient has small cell neuroendocrine carcinoma of the cervix (NECC); **AND**

- Used as first-line or subsequent therapy (if not used previously as first-line therapy) for persistent, recurrent, or metastatic disease; **AND**
- Used in combination with etoposide **AND** either cisplatin or carboplatin

§ Genomic Aberration/Mutational Driver Targeted Therapies (Note: not all inclusive, refer to guidelines for appropriate use)				
Sensitizing <i>EGFR</i> mutation-positive tumors	<i>ALK</i> rearrangement-positive tumors	<i>ROS1</i> rearrangement-positive tumors	<i>BRAF</i> V600E-mutation positive tumors	<i>NTRK1/2/3</i> gene fusion positive tumors
– Afatinib – Erlotinib – Dacomitinib – Gefitinib – Osimertinib – Amivantamab (<i>exon-20 insertion</i>) – Mobocertinib (<i>exon-20 insertion</i>)	– Alectinib – Brigatinib – Ceritinib – Crizotinib – Lorlatinib	– Ceritinib – Crizotinib – Entrectinib – Lorlatinib	– Dabrafenib ± trametinib – Vemurafenib	– Larotrectinib – Entrectinib
PD-L1 tumor expression ≥ 1%	<i>MET</i> exon-14 skipping mutations	<i>RET</i> rearrangement-positive tumors	<i>KRAS</i> G12C mutation positive tumors	<i>ERBB2 (HER2)</i> mutation positive tumors
– Pembrolizumab – Atezolizumab – Nivolumab + ipilimumab – Cemiplimab – Tremelimumab + durvalumab	– Capmatinib – Crizotinib – Tepotinib	– Selpercatinib – Cabozantinib – Pralsetinib	– Sotorasib – Adagrasib	– Fam-trastuzumab deruxtecan-nxki – Ado-trastuzumab emtansine

Δ Notes:

- Patients responding to therapy who relapse ≥ 6 months after discontinuation due to duration (i.e., receipt of 24 months of therapy) are eligible to re-initiate PD-directed therapy.
- Patients who complete adjuvant therapy and progress ≥ 6 months after discontinuation are eligible to re-initiate PD-directed therapy for metastatic disease.
- Patients whose tumors, upon re-biopsy, demonstrate a change in actionable mutation (e.g., MSS initial biopsy; MSI-H subsequent biopsy) may be eligible to re-initiate PD-directed therapy and will be evaluated on a case-by-case basis.

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Ⓢ Orphan Drug

IV. Renewal Criteria ^{Δ 1,3}

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe or life-threatening infusion-related reactions, immune-mediated adverse reactions

(e.g., pneumonitis, hepatitis, colitis, endocrinopathies, nephritis with renal dysfunction, dermatology reactions, pancreatitis, etc.), complications of allogeneic hematopoietic stem cell transplantation (HCST), etc.: **AND**

NSCLC (single-agent use as consolidation therapy)

- Patient has not exceeded a maximum of 12 months of therapy

Continuation Maintenance Therapy for NSCLC

- *Refer to Section III for criteria*

Hepatobiliary Cancers

- Cases for patients with HCC who use treatment as part of STRIDE and experience disease progression but who are clinically stable and still deriving clinical benefit will be reviewed on a case-by-case basis.

Continuation Maintenance Therapy for SCLC

- *Refer to Section III for criteria*

V. Dosage/Administration ^{Δ 1,7,8,12,17}

Indication	Dose
Non-Small Cell Lung Cancer (NSCLC)	<u>Single agent:</u> • Weight ≥ 30 kg: Administer 10 mg/kg intravenously every 14 days OR 1,500 mg intravenously every 28 days until disease progression or unacceptable toxicity • Weight < 30 kg: Administer 10 mg/kg intravenously every 14 days until disease progression or unacceptable toxicity • <i>NOTE:</i> <i>Use as consolidation therapy for unresectable stage II-III disease may continue up to a maximum of 12 months in patients without disease progression or unacceptable toxicity.</i> <u>In combination with Tremelimumab-actl* and Platinum-Based Chemotherapy§:</u> • Weight ≥ 30 kg: Administer 1,500 mg intravenously every 21 days x 4 cycles, followed by a maintenance dose of 1,500 mg every 28 days thereafter, until disease progression or unacceptable toxicity • Weight < 30 kg: Administer 20 mg/kg intravenously every 21 days x 4 cycles, followed by a maintenance dose of 20 mg/kg every 28 days thereafter, until disease progression or unacceptable toxicity <i>*Note:</i> <i>Refer to the Prescribing Information for tremelimumab-actl dosing information</i> <i>§</i> <i>If patients receive fewer than 4 cycles of platinum-based chemotherapy, the remaining cycles of Tremelimumab-actl (up to a total of 5) should be given</i>

	<i>after the platinum-based chemotherapy phase, in combination with IMFINZI, every 4 weeks.</i>
Small Cell Lung Cancer (SCLC)	<u>Weight ≥30 kg:</u> Administer 1,500 mg intravenously in combination with chemotherapy every 21 days x 4 cycles*, followed by a maintenance dose of 1,500 mg as a single agent every 28 days thereafter, until disease progression or unacceptable toxicity <u>Weight <30 kg:</u> Administer 20 mg/kg intravenously in combination with chemotherapy every 21 days x 4 cycles*, followed by a maintenance dose of 10 mg/kg as a single agent every 14 days thereafter, until disease progression or unacceptable toxicity *Note: <i>Patients may receive up to 2 additional cycles in combination with chemotherapy based on response and tolerability after the initial 4 cycles (6 cycles of combination therapy in total) ^s</i>
Hepatocellular Carcinoma	<u>Single agent:</u> Administer 1,500 mg intravenously every 4 weeks until disease progression or unacceptable toxicity <u>STRIDE (Single Tremelimumab Regular Interval Durvalumab):</u> • Weight ≥30 kg: Administer 1,500 mg intravenously following a single dose of tremelimumab-actl* at Day 1 of Cycle 1, followed by a maintenance dose of 1,500 mg as a single agent every 28 days thereafter, until disease progression or unacceptable toxicity • Weight <30 kg: Administer 20 mg/kg intravenously following a single dose of tremelimumab-actl* at Day 1 of Cycle 1, followed by a maintenance dose of 20 mg/kg as a single agent every 28 days thereafter, until disease progression or unacceptable toxicity *Note: <i>Refer to the Prescribing Information for tremelimumab-actl dosing information</i>
Biliary Tract Cancer	<u>Weight ≥30 kg:</u> Administer 1,500 mg intravenously in combination with chemotherapy every 21 days for up to 8 cycles, followed by a maintenance dose of 1,500 mg as a single agent every 28 days thereafter, until disease progression or unacceptable toxicity <u>Weight <30 kg:</u> Administer 20 mg/kg intravenously in combination with chemotherapy every 21 days for up to 8 cycles, followed by a maintenance dose of 20 mg/kg as a single agent every 28 days thereafter, until disease progression or unacceptable toxicity
Cervical Cancer	<u>Weight ≥30 kg:</u> Administer 1,500 mg intravenously in combination with chemotherapy every 21 days x 4 cycles, followed by a maintenance dose of 1,500 mg as a single

	agent every 28 days thereafter, until disease progression or unacceptable toxicity <u>Weight <30 kg:</u> Administer 20 mg/kg intravenously in combination with chemotherapy every 21 days x 4 cycles, followed by a maintenance dose of 10 mg/kg as a single agent every 14 days thereafter, until disease progression or unacceptable toxicity
<u>Dosing should be calculated using actual body weight and not flat dosing (as applicable) based on the following:</u> • Patient weight > 30 kg and <75 kg: Use 20 mg/kg dosing <i>Note: This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide. Patient-specific variables should be taken into account.</i>	

VI. Billing Code/Availability Information

HCPCS Code:

- J9173 – Injection, durvalumab, 10 mg; 1 billable unit = 10 mg

NDC(s):

- Imfinzi 120 mg/2.4 mL single-dose vial: 00310-4500-xx
- Imfinzi 500 mg/10 mL single-dose vial: 00310-4611-xx

VII. References

1. Imfinzi [package insert]. Wilmington, DE; AstraZeneca Pharmaceuticals LP; November 2022. Accessed January 2023.
2. Massard C, Gordon MS, Sharma S, et al. Safety and Efficacy of Durvalumab (MEDI4736), an Anti-Programmed Cell Death Ligand-1 Immune Checkpoint Inhibitor, in Patients With Advanced Urothelial Bladder Cancer. J Clin Oncol. 2016 Sep 10;34(26):3119-25.
3. Referenced with permission from the NCCN Drugs and Biologics Compendium (NCCN Compendium®) durvalumab. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®, NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed January 2023.
4. Antonia SJ, Villegas A, Daniel D, et al. Durvalumab after Chemoradiotherapy in Stage III Non-Small-Cell Lung Cancer. N Engl J Med. 2017 Sep 8.
5. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Non-Small Cell Lung Cancer. Version 1.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN

Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed January 2023.

6. Gupta S, Sonpavde G, Grivas P, et al. Defining “platinum-ineligible” patients with metastatic urothelial cancer (mUC). *J Clin Oncol*. 2019 Mar 1;37(7_suppl):451.
7. Paz-Ares L, Dvorkin M, Chen Y, et al. Durvalumab plus platinum-etoposide versus platinum-etoposide in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial. *Lancet*. 2019 Nov 23;394(10212):1929-1939. Doi: 10.1016/S0140-6736(19)32222-6. Epub 2019 Oct 4.
8. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Small Cell Lung Cancer. Version 3.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed January 2023.
9. Powles T, O'Donnell PH, Massard C, et al. Efficacy and Safety of Durvalumab in Locally Advanced or Metastatic Urothelial Carcinoma: Updated Results From a Phase ½ Open-label Study. *JAMA Oncol*. 2017 Sep 14;3(9):e172411. Doi: 10.1001/jamaoncol.2017.2411. Epub 2017 Sep 14.
10. Goldman JW, Dvorkin M, Chen Y, et al. Durvalumab, with or without tremelimumab, plus platinum-etoposide versus platinum-etoposide alone in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): updated results from a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol*. 2021 Jan;22(1):51-65. Doi: 10.1016/S1470-2045(20)30539-8.
11. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Hepatobiliary Cancers. Version 5.2022. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed January 2023.
12. Abou-Alfa GK, Chan SL, Kudo M, et al. Phase 3 randomized, open-label, multicenter study of tremelimumab (T) and durvalumab (D) as first-line therapy in patients (pts) with unresectable hepatocellular carcinoma (uHCC): HIMALAYA. *Journal of Clinical Oncology* 2022 40:4_suppl, 379-379.
13. Govindan R, Aggarwal C, Antonia SJ, et al. Society for Immunotherapy of Cancer (SITC) clinical practice guideline on immunotherapy for the treatment of lung cancer and mesothelioma. *Journal for ImmunoTherapy of Cancer* 2022;10:e003956. Doi: 10.1136/jitc-2021-003956.

14. Oh DY, He AR, Qin S, et al. A phase 3 randomized, double-blind, placebo-controlled study of durvalumab in combination with gemcitabine plus cisplatin (GemCis) in patients (pts) with advanced biliary tract cancer (BTC): TOPAZ-1. J Clin Oncol. 2022 Feb 1;40(4_suppl):378-378.
15. Abou-Alfa GK, Chan SL, Furuse J, et al. A randomized, multicenter phase 3 study of durvalumab (D) and tremelimumab (T) as first-line treatment in patients with unresectable hepatocellular carcinoma (HCC): HIMALAYA study. Journal of Clinical Oncology 36, no. 15_suppl. DOI: 10.1200/JCO.2018.36.15_suppl.TPS4144.
16. Johnson ML, Cho BC, Luft A, et al; POSEIDON investigators. Durvalumab With or Without Tremelimumab in Combination With Chemotherapy as First-Line Therapy for Metastatic Non-Small-Cell Lung Cancer: The Phase III POSEIDON Study. J Clin Oncol. 2022 Nov 3;JCO2200975. doi: 10.1200/JCO.22.00975.
17. Paz-Ares L, Dvorkin M, Chen Y, et al; CASPIAN investigators. Durvalumab plus platinum-etoposide versus platinum-etoposide in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial. Lancet. 2019 Nov 23;394(10212):1929-1939. doi: 10.1016/S0140-6736(19)32222-6.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C22.0	Liver cell carcinoma
C22.1	Intrahepatic bile duct carcinoma
C22.8	Malignant neoplasm of liver, primary, unspecified as to type
C22.9	Malignant neoplasm of liver, not specified as primary or secondary
C23	Malignant neoplasm of gallbladder
C24.0	Malignant neoplasm of other and unspecified parts of biliary tract
C24.8	Malignant neoplasm of overlapping sites of biliary tract
C24.9	Malignant neoplasm of biliary tract, unspecified
C33	Malignant neoplasm of trachea
C34.00	Malignant neoplasm of unspecified main bronchus
C34.01	Malignant neoplasm of right main bronchus
C34.02	Malignant neoplasm of left main bronchus
C34.10	Malignant neoplasm of upper lobe, unspecified bronchus or lung
C34.11	Malignant neoplasm of upper lobe, right bronchus or lung
C34.12	Malignant neoplasm of upper lobe, left bronchus or lung
C34.2	Malignant neoplasm of middle lobe, bronchus or lung
C34.30	Malignant neoplasm of lower lobe, unspecified bronchus or lung
C34.31	Malignant neoplasm of lower lobe, right bronchus or lung
C34.32	Malignant neoplasm of lower lobe, left bronchus or lung
C34.80	Malignant neoplasm of overlapping sites of unspecified bronchus and lung
C34.81	Malignant neoplasm of overlapping sites of right bronchus and lung

IMFINZI® (durvalumab) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2023, Magellan Rx Management

ICD-10	ICD-10 Description
C34.82	Malignant neoplasm of overlapping sites of left bronchus and lung
C34.90	Malignant neoplasm of unspecified part of unspecified bronchus or lung
C34.91	Malignant neoplasm of unspecified part of right bronchus or lung
C34.92	Malignant neoplasm of unspecified part of left bronchus or lung
C53.0	Malignant neoplasm of endocervix
C53.1	Malignant neoplasm of exocervix
C53.8	Malignant neoplasm of overlapping sites of cervix uteri
C53.9	Malignant neoplasm of cervix uteri, unspecified
C7A.1	Malignant poorly differentiated neuroendocrine tumors
C78.00	Secondary malignant neoplasm of unspecified lung
C78.01	Secondary malignant neoplasm of right lung
C78.02	Secondary malignant neoplasm of left lung
C79.31	Secondary malignant neoplasm of brain
C79.51	Secondary malignant neoplasm of bone
C79.52	Secondary malignant neoplasm of bone marrow
Z85.118	Personal history of other malignant neoplasm of bronchus and lung

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC

Nondiscrimination & Language Access Policy

Aspirus Health Plan, Inc. complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, sex, sexual orientation, or gender identity. We do not exclude people or treat them differently because of race, color, national origin, age, disability, sex, sexual orientation, or gender identity.

We will:

Provide free aids and services to people with disabilities to communicate effectively with us, such as:

- Qualified sign language interpreters
- Written information in other formats (large print, audio, accessible electronic formats, other formats)

Provide free language services to people whose primary language is not English, such as:

- Qualified interpreters
- Information written in other languages

If you need these services, contact us at the phone number shown on the inside cover of this contract, your id card, or aspirushealthplan.com.

If you believe that we have failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, sex, sexual orientation, or gender identity, you can file a grievance with:

Nondiscrimination Grievance Coordinator
Aspirus Health Plan, Inc.
PO Box 1062
Minneapolis, MN 55440
Phone: 1.866.631.5404 (TTY: 711)
Fax: 763.847.4010
Email: customerservice@aspirushealthplan.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Nondiscrimination Grievance Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at:

U.S. Department of Health and Human Services
200 Independence Avenue, SW
Room 509F, HHH Building
Washington, D.C. 20201
1-800-368-1019, 800-537-7697 (TDD)

Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

Language Assistance Services

Albanian: KUJDES: Nëse flitni shqip, për ju ka në dispozicion shërbime të asistencës gjuhësore, pa pagesë. Telefononi në 1.866.631.5404 (TTY: 711).

Arabic: تنبيه: إذا كنت تتحدث اللغة العربية، فإن خدمات المساعدة اللغوية متاحة لك مجاناً. اتصل بن اعلی رقم الهاتف 1.866.631.5404 (رقم هاتف الصم والبك : 711)

French: ATTENTION : Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelezle 1.866.631.5404 (ATS : 711).

German: ACHTUNG: Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zur Verfügung. Rufnummer: 1.866.631.5404 (TTY: 711).

Hindi: _यान द_ : य_द आप िहंदी बोलते ह_ तो आपके िलए मु_त म_ भाषा सहायता सेवाएं उपल_ध ह_। 1.866.631.5404 (TTY: 711) पर कॉल कर_।

Hmong: LUS CEEV: Yog tias koj hais lus Hmoob, cov kev pab txog lus, muaj kev pab dawb rau koj. Hu rau 1.866.631.5404 (TTY: 711).

Korean: 주의: 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 1.866.631.5404 (TTY: 711)번으로 전화해 주십시오.

Polish: UWAGA: Jeżeli mówisz po polsku, możesz skorzystać z bezpłatnej pomocy językowej. Zadzwoń pod numer 1.866.631.5404 (TTY: 711).

Russian: ВНИМАНИЕ: Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода. Звоните 1.866.631.5404 (телетайп: 711).

Spanish: ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1.866.631.5404 (TTY: 711).

Tagalog: PAUNAWA: Kung nagsasalita ka ng Tagalog, maaari kang gumamit ng mga serbisyo ng tulong sa wika nangwalang bayad. Tumawag sa 1.866.631.5404 (TTY: 711)

Traditional Chinese: 注意: 如果您使用繁體中文, 您可以免費獲得語言援助服務。請致電 1.866.631.5404 (TTY: 711)。

Vietnamese: CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 1.866.631.5404 (TTY: 711).

Pennsylvania Dutch: Wann du Deitsch (Pennsylvania German / Dutch) schwetzscht, kannscht du mitaus Koschte ebbergricke, ass dihr helft mit die englisch Schprooch. Ruf selli Nummer uff: Call 1.866.631.5404 (TTY: 711).

Lao: ໄປດຊາຍ: ຖ້າວ່າ ທ່ານເວົ້າພາສາ ລາວ, ການບໍລິການຊ່ວຍເຫຼືອດ້ານພາສາ, ໂດຍບໍ່ເສັຽຄ່າ, ແມ່ນມີພ້ອມໃຫ້ທ່ານ. ໂທ 1.866.631.5404 (TTY: 711).