Hemophilia Products – von Willebrand Factor: Vonvendi®
(Intravenous)

Document Number: IC-0344

Last Review Date: 06/01/2023
Date of Origin: 12/16/2014
Dates Reviewed: 12/2014, 04/2015, 05/2015, 09/2015, 12/2015, 03/2016, 06/2016, 12/2016, 06/2017, 09/2017, 11/2017, 11/2018, 03/2019, 10/2019, 02/2020, 06/2021, 03/2022, 06/2022, 06/2023

I. Length of Authorization

Coverage is provided for 3 months and may be renewed thereafter, unless otherwise specified*.

Note: The cumulative amount of medication the patient has on-hand will be taken into account for authorizations. Up to 5 'on-hand' doses for the treatment of acute bleeding episodes will be permitted at the time of the authorization request.

** Initial and renewal authorization periods may vary by specific covered indication*

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Vonvendi 450-850 units: 82 vials per 90-day supply
- Vonvendi 900-1700 units: 41 vials per 90-day supply

B. Max Units (per dose and over time) [HCPCS Unit]:

- 36,800 billable units per 90 day supply

III. Initial Approval Criteria [1-3]

Hemophilia Management Program
Requirements for inhibitor tests are a part of the hemophilia management program. This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide.

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

Von Willebrand Disease (vWD) † Φ

Publication of this policy on this website is an authorized use of proprietary and copyrighted information created in collaboration with Magellan Rx Management. This policy may not be reproduced or distributed without the express written permission of Magellan Rx Management.
© 2024 Magellan Rx Management

- Diagnosis of von Willebrand disease has been confirmed by blood coagulation and von Willebrand factor testing; **AND**
  - Used for the treatment of spontaneous and trauma-induced bleeding episodes in at least one of the following:
    - Patient has severe vWD
    - Patient has mild or moderate vWD and the use of desmopressin is known or suspected to be ineffective or contraindicated
    - Used for perioperative management of bleeding *(Note: Authorizations valid for 1 month)*; **OR**
  - Used for routine prophylaxis to reduce the frequency of bleeding episodes; **AND**
    - Patient has severe Type 3 vWD and is receiving on-demand therapy

Hemophilia Management Program
For minimally treated patients (< 50 exposure days to factor products) previously receiving a different factor product, inhibitor testing is required at baseline, then at every comprehensive care visit (yearly for the mild and moderate patients, semi-annually for the severe patients)

† FDA Approved Indication(s); ‡ Compendia recommended Indication(s); Φ Orphan Drug

IV. Dispensing Requirements for Rendering Providers (Hemophilia Management Program)

- Prescriptions cannot be filled without an expressed need from the patient, caregiver or prescribing practitioner. Auto-filling is not allowed.
- Monthly, rendering provider must submit for authorization of dispensing quantity before delivering factor product. Information submitted must include:
  - Original prescription information, requested amount to be dispensed, vial sizes available to be ordered from the manufacturer, and patient clinical history (including patient product inventory and bleed history)
  - Factor dose should not exceed +1% of the prescribed dose and a maximum of three vials may be dispensed per dose. If unable to provide factor dosing within the required threshold, below the required threshold, the lowest possible dose able to be achieved above +1% should be dispensed. Prescribed dose should not be increased to meet assay management requirements.
- The cumulative amount of medication(s) the patient has on-hand should be taken into account when dispensing factor product. Patients should not have more than 5 extra doses on-hand for the treatment of acute bleeding episodes.
- Dispensing requirements for renderings providers are a part of the hemophilia management program. This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
©2023, Magellan Rx Management

V. Renewal Criteria [1-3]

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the indication-specific relevant criteria identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: symptoms of allergic-anaphylactic reactions (anaphylaxis, dyspnea, rash, etc.), thromboembolic events (thromboembolism, pulmonary embolism), development of neutralizing antibodies (inhibitors), etc.; **AND**
- Any increases in dose must be supported by an acceptable clinical rationale (i.e. weight gain, half-life study results, increase in breakthrough bleeding when patient is fully adherent to therapy, etc.); **AND**
- The cumulative amount of medication(s) the patient has on-hand will be taken into account when authorizing. The authorization will allow up to 5 doses on-hand for the treatment of acute bleeding episodes as needed for the duration of the authorization; **AND**

Treatment of spontaneous and trauma-induced bleeding episodes

- Renewals will be approved for a 6 month authorization period

Perioperative management of surgical bleeding

- Coverage may NOT be renewed

Routine prophylaxis to reduce the frequency of bleeding episodes

- Renewals will be approved for a 12 month authorization period; **AND**
- Patient has demonstrated a beneficial response to therapy (i.e., the frequency of bleeding episodes has decreased from pre-treatment baseline)

VI. Dosage/Administration [1-3]

Indication	Dose
Treatment and control of bleeding episodes VWD	• For each bleeding episode, administer the first dose of Vonvendi with an approved recombinant (non-von Willebrand factor containing) factor VIII if factor VIII baseline levels are below 40% or are unknown. • If recombinant factor VIII is required, give recombinant factor VIII within 10 minutes of completing Vonvendi infusion at a ratio of 1.3:1 (i.e., 30% more Vonvendi than recombinant factor VIII, based on the approximate mean recoveries of 1.5 and 2 IU/dL for Vonvendi and recombinant factor VIII, respectively). Minor: Loading dose: 40-50 IU/kg Maintenance dose: 40-50 IU/kg every 8-24 hours as clinically required Major:

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
©2023, Magellan Rx Management

Indication	Dose
	Loading dose: 50-80 IU/kg Maintenance dose: 40-60 IU/kg every 8-24 hours for approximately 2 to 3 days as clinically required
Perioperative management of bleeding VWD	<u>Elective Surgical Procedure</u> A preoperative dose may be administered 12-24 hours prior to surgery to allow the endogenous factor VIII levels to increase to at least 30 IU/dL (minor surgery) or 60 IU/dL (major surgery) before the loading dose (1 hour preoperative dose) of rVWF, with or without recombinant factor VIII, is administered. • Ensure baseline FVIII:C level is available prior to determining the need for 12-24 hr preoperative dose. FVIII:C level should also be assessed within 3 hours prior to initiating the surgical procedure. If the level is at the recommended minimum target levels (30 IU/dL for minor surgery and 60 IU/dL for major surgery), administer a dose of Vonvendi alone (without factor VIII treatment) within 1 hour prior to the procedure. If the FVIII:C level is below the recommended minimum target level, administer complete dose of Vonvendi followed by recombinant factor VIII within 10 minutes to raise VWF:RCo and FVIII:C. • Assess baseline VWF:RCo levels within 3 hours of administration of the 12-24 hr preoperative dose. If the 12-24 hour preoperative dose is not administered, then assess baseline level VWF:RCo prior to surgery. When possible, measure incremental recovery (IR) for Vonvendi before surgery. For calculation of IR, measure baseline plasma VWF:RCo. Then infuse a dose of 50 IU/kg of Vonvendi. Measure VWF:RCo, 30 minutes after infusion of Vonvendi. • Use the following formula to calculate IR: IR= [Plasma VWF:RCo at 30 minutes (IU/dL) – Plasma VWF:RCo at baseline (IU/dL)]/Dose(IU/kg). <u>Emergency Surgical Procedure</u> • A 12-24 hr preoperative dose may not be feasible in subjects requiring emergency surgery. Baseline VWF:RCo and FVIII:C levels should be assessed within 3 hours prior to initiating the surgical procedure if it is feasible. The loading dose (1 hour preoperative dose) can be calculated as the difference in the target peak and baseline plasma VWF:RCo levels divided by the IR. If the IR is not available, assume an IR of 2.0 IU/dL per IU/kg. If baseline VWF:RCo and FVIII:C is not available, as a general guidance a loading dose (1 hour preoperative dose), 40 to 60 IU/kg VWF:RCo, should be administered. Additionally, recombinant factor VIII at a dose of 30 to 45 IU/kg may be infused sequentially, preferably within 10 minutes after the Vonvendi infusion in patients whose factor VIII plasma levels already are (or are highly likely to be) less than 40 to 50 IU/dL for minor surgery or 80 to 100 IU/dL for major surgery.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
©2023, Magellan Rx Management

Indication	Dose
	Note: refer to the package insert for recommended VWF:RCo and FVIII:C target peak plasma levels and dosing guidelines for perioperative management of bleeding.
Routine Prophylaxis to Reduce the Frequency of Bleeding Episodes	• For initiation of prophylactic treatment, administer 40 to 60 IU/kg twice weekly. – Adjust prophylaxis dose up to 60 IU/kg twice weekly if breakthrough bleeding* occurs in joints or if severe bleeding occurs. **Treat breakthrough bleeding as per the dosing guidelines for minor and major bleeding.*

VII. Billing Code/Availability Information

HCPCS Code & NDC(s):

Drug	Manufacturer	HCPCS Code	1 Billable Unit Equiv.	Vial Size	NDC
Vonvendi	Baxalta US Inc.	J7179	1 IU	450-850 IU	00944-7551-xx
				900-1700 IU	00944-7553-xx

VIII. References

1. Vonvendi [package insert]. Lexington, MA; Takeda Phamaceuticals USA, Inc.; March 2023. Accessed May 2023.
2. MASAC Recommendations Concerning Products Licensed For The Treatment Of Hemophilia And Other Bleeding Disorders. Revised April 2022 National Hemophilia Foundation. MASAC Document #272; April 2022. Available at: http://www.hemophilia.org. Accessed May 2023.
3. Guidelines for the Management of Hemophilia. 3rd Edition. World Federation of Hemophilia 2020. Available at: https://www1.wfh.org/publications/files/pdf-1863.pdf. Accessed May 2023.
4. Graham A1, Jaworski K. Pharmacokinetic analysis of anti-hemophilic factor in the obese patient. Haemophilia. 2014 Mar;20(2):226-9.
5. Croteau SE1, Neufeld EJ. Transition considerations for extended half-life factor products. Haemophilia. 2015 May;21(3):285-8.
6. Mingot-Castellano, et al. Application of Pharmacokinetics Programs in Optimization of Haemostatic Treatment in Severe Hemophilia a Patients: Changes in Consumption, Clinical Outcomes and Quality of Life. Blood. 2014 December; 124 (21).
7. MASAC Recommendation Concerning Prophylaxis. 2022 National Hemophilia Foundation. MASAC Document #267; April 2022. Available at: http://www.hemophilia.org. Accessed May 2023.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
©2023, Magellan Rx Management

8. MASAC Recommendations Regarding the Treatment of von Willebrand Disease. MASAC Document #266; March 2021. Available at: 266.pdf (hemophilia.org). Accessed May 2023.
9. Connell NT, Flood VH, Brignardello-Petersen R, et al. ASH ISTH NHF WFH 2021 guidelines on the management of von Willebrand disease. Blood Advances, Volume 5, Issue 1, 2021, Pages 301-325, https://doi.org/10.1182/bloodadvances.2020003264.
10. Franchini M, Seidizadeh, Mannucci P, et al. Prophylactic management of patients with von Willebrand disease. Ther Adv Hematol. 2021; 12: 20406207211064064. Published online 2021 Dec 22. doi: 10.1177/20406207211064064
11. First Coast Service Options, Inc. Local Coverage Article: Billing and Coding: Hemophilia Clotting Factors (A56482). Centers for Medicare & Medicaid Services Inc. Updated on 10/28/2022 with effective date 10/01/2022. Accessed May 2023.
12. Palmetto GBA. Local Coverage Article: Billing and Coding: Guidance for Anti-Inhibitor Coagulant Complex (AICC) National Coverage Determination (NCD) 110.3 (A56065). Centers for Medicare & Medicaid Services Inc. Updated on 11/14/2022 with effective date 11/24/2022. Accessed May 2023.
13. Novitas Solutions, Inc. Local Coverage Article: Billing and Coding: Hemophilia Factor Products (A56433). Centers for Medicare & Medicaid Services Inc. Updated on 10/14/2022 with effective date 10/01/2022. Accessed May 2023.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
D68.01	Von Willebrand disease, type 1
D68.020	Von Willebrand disease, type 2A
D68.021	Von Willebrand disease, type 2B
D68.022	Von Willebrand disease, type 2M
D68.023	Von Willebrand disease, type 2N
D68.03	Von Willebrand disease, type 3
D68.04	Acquired von Willebrand disease
D68.09	Other von Willebrand disease

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Articles (LCAs) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: https://www.cms.gov/medicare-coverage-database/search.aspx. Additional indications may be covered at the discretion of the health plan.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
©2023, Magellan Rx Management

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCA/LCD):

Jurisdiction(s): N	**NCD/LCD Document (s):** A56482
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a56482&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	

Jurisdiction(s): J,M	**NCD/LCD Document (s):** A56065
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a56065&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	

Jurisdiction(s): H,L	**NCD/LCD Document (s):** A56433
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a56433&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	**Applicable State/US Territory**	**Contractor**
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
©2023, Magellan Rx Management

Nondiscrimination & Language Access Policy

Aspirus Health Plan, Inc. complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, sex, sexual orientation, or gender identity. *We* do not exclude people or treat them differently because of race, color, national origin, age, disability, sex, sexual orientation, or gender identity.

We will:
Provide free aids and services to people with disabilities to communicate effectively with us, such as:
- Qualified sign language interpreters
- Written information in other formats (large print, audio, accessible electronic formats, other formats)

Provide free language services to people whose primary language is not English, such as:
- Qualified interpreters
- Information written in other languages

If *you* need these services, contact *us* at the phone number shown on the inside cover of this *contract*, *your* id card, or aspirushealthplan.com.

If *you* believe that *we* have failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, sex, sexual orientation, or gender identity, *you* can file a grievance with:

Nondiscrimination Grievance Coordinator
Aspirus Health Plan, Inc.
PO Box 1062
Minneapolis, MN 55440
Phone: 1.866.631.5404 (TTY: 711)
Fax: 763.847.4010
Email: customerservice@aspirushealthplan.com

You can file a grievance in person or by mail, fax, or email. If *you* need help filing a grievance, the Nondiscrimination Grievance Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at https://ocrportal.hhs.gov/ocr/portal/lobby.jsf, or by mail or phone at:

U.S. Department of Health and Human Services
200 Independence Avenue, SW
Room 509F, HHH Building
Washington, D.C. 20201
1-800-368-1019, 800-537-7697 (TDD)

Complaint forms are available at http://www.hhs.gov/ocr/office/file/index.html.

Language Assistance Services

Albanian: KUJDES: Nëse flitni shqip, për ju ka në dispozicion shërbime të asistencës gjuhësore, pa pagesë.Telefononi në 1.866.631.5404 (TTY: 711).

Arabic تنبيه: إذا كنت تتحدث اللغة العربية، فإن خدمات المساعدة اللغوية متاحة لك مجاناً . اتصل بن اعلى رقم الهاتف 1.866.631.5404(رقم هاتف الصم والبك : 711)

French: ATTENTION : Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelezle 1.866.631.5404 (ATS : 711).

German: ACHTUNG: Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zurVerfügung. Rufnummer: 1.866.631.5404 (TTY: 711).

Hindi: _यान द_: य_द आप िहंदी बोलते ह_ तो आपके िलए मु_त म_ भाषा सहायता सेवाएं उपल_ध ह_। 1.866.631.5404 (TTY: 711) पर कॉल कर_।

Hmong: LUS CEEV: Yog tias koj hais lus Hmoob, cov kev pab txog lus, muaj kev pab dawb rau koj. Hu rau 1.866.631.5404 (TTY: 711).

Korean: 주의: 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 1.866.631.5404 (TTY: 711)번으로 전화해 주십시오.

Polish: UWAGA: Jeżeli mówisz po polsku, możesz skorzystać z bezpłatnej pomocy językowej. Zadzwoń pod numer 1.866.631.5404 (TTY: 711).

Russian: ВНИМАНИЕ: Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода.Звоните 1.866.631.5404 (телетайп: 711).

Spanish: ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1.866.631.5404 (TTY: 711).

Tagalog: PAUNAWA: Kung nagsasalita ka ng Tagalog, maaari kang gumamit ng mga serbisyo ng tulong sa wika nangwalang bayad. Tumawag sa 1.866.631.5404 (TTY: 711)

Traditional Chinese: 注意：如果您使用繁體中文，您可以免費獲得語言援助服務。請 致電 1.866.631.5404 (TTY:711).

Vietnamese: CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 1.866.631.5404 (TTY: 711).

Pennsylvania Dutch: Wann du Deitsch (Pennsylvania German / Dutch) schwetzscht, kannscht du mitaus Koschte ebbergricke, ass dihr helft mit die englisch Schprooch. Ruf selli Nummer uff: Call 1.866.631.5404 (TTY: 711).

Lao: ໂປດຊາບ: ຖ້າວ່າ ທ່ານເວົ້າພາສາ ລາວ, ການບໍລິການຊ່ວຍເຫຼືອດ້ານພາສາ,ໂດຍບໍ່ເສັ້ງຄ່າ, ແມ່ນມີພ້ອມໃຫ້ທ່ານ. ໂທຣ 1.866.631.5404 (TTY:711).

60017 12-22 ©2022 Aspirus Health Plan, Inc. All rights reserved.