

Lutathera® (lutetium Lu 177 dotatate) (Intravenous)

Document Number: IC-0352

Last Review Date: 07/05/2023

Date of Origin: 02/06/2018

Dates Reviewed: 02/2018, 07/2018, 07/2019, 07/2020, 07/2021, 07/2022, 07/2023

I. Length of Authorization

Coverage will be provided for 1 year (4 doses only) and may NOT be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- N/A

B. Max Units (per dose and over time) [HCPCS Unit]:

- 200 billable units (7.4 GBq = 200 mCi) every 8 weeks for a total of 4 doses

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**
- Patient has a negative pregnancy test (in females) prior to initiating treatment; **AND**
- Patient's disease is somatostatin receptor-positive in all tumor lesions (OctreoScan uptake $\geq$ normal liver); **AND**
- Patient has well-differentiated disease with a Ki67 labeling index score of $\leq 20\%$ (*NOTE: Excluding use in Well-differentiated grade 3 NET and Pheochromocytoma or Paraganglioma*); **AND**
- Patient has not received any long-acting somatostatin analogues (e.g., octreotide LAR, pasireotide LAR, lanreotide depot, etc.) within the previous 4 weeks OR short-acting somatostatin analogues (e.g., octreotide, pasireotide, etc.) within 24 hours prior to therapy; **AND**
- Will be used in combination with a long-acting somatostatin analog (e.g., octreotide LAR, lanreotide depot, etc.) given as a single-injection (between 4-24 hours) following each Lutathera infusion; **AND**

Neuroendocrine and Adrenal Tumors ^{1,3,4}

- Used for Gastroenteropancreatic Neuroendocrine Tumors (GEP-NETs) † Φ ; **AND**

- Patient has progressive locally advanced or metastatic disease; **AND**
 - Disease has progressed on long-acting octreotide or lanreotide; **OR**
 - Used for well-differentiated grade 3 NET ‡; **AND**
 - Patient has unresectable locally advanced or metastatic disease; **AND**
 - Patient has favorable biology (e.g., relatively low Ki-67 [<55%], positive SSTR-based PET imaging) that has clinically significant tumor burden or evidence of disease progression; **OR**
 - Used for bronchopulmonary/thymic disease ‡; **AND**
 - Disease has progressed on long-acting octreotide or lanreotide; **AND**
 - Used as subsequent therapy for recurrent and/or unresectable disease; **OR**
 - Used as primary therapy OR subsequent therapy (as alternate primary therapy) if progression on primary therapy; **AND**
 - Patient has recurrent and/or metastatic disease with clinically significant tumor burden and low grade (typical carcinoid) histology, evidence of disease progression, intermediate grade (atypical carcinoid) histology, or symptomatic disease; **OR**
 - Used for gastrointestinal disease ‡; **AND**
 - Disease has progressed on long-acting octreotide or lanreotide; **AND**
 - Patient has recurrent, locoregional advanced and/or distant metastatic disease; **AND**
 - Used as subsequent therapy OR as alternative-front-line therapy for clinically significant tumor burden; **OR**
 - Patient has poorly controlled carcinoid syndrome ‡; **AND**
 - Disease has progressed on long-acting octreotide or lanreotide; **AND**
 - Patient has persistent symptoms (i.e., flushing, diarrhea); **OR**
 - Used for pancreatic disease ‡; **AND**
 - Disease has progressed on long-acting octreotide or lanreotide; **AND**
 - Patient has symptomatic, clinically significant tumor burden OR progressive recurrent, locoregional advanced, and/or distant metastatic disease; **AND**
 - Used as subsequent therapy OR as alternative-front-line therapy; **OR**
 - Used for Pheochromocytoma or Paraganglioma ‡; **AND**
 - Used as primary treatment for locally unresectable or metastatic disease
- † FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Ⓢ Orphan Drug

IV. Renewal Criteria

Coverage cannot be renewed.

V. Dosage/Administration¹

Indication	Dose
All Indications	- Administer 7.4 GBq (200 mCi) every 8 weeks ($\pm$ 1 week) for a total of 4 doses. Administer a single dose of long-acting somatostatin analog between 4 to 24 hours after each Lutathera dose. (Long-acting somatostatin analog may not be repeated until after the next scheduled dose of Lutathera to provide the 4-week drug-free interval. Short-acting octreotide may be administered up to 24 hours prior to each Lutathera dose) - Initiate recommended intravenous amino acid solution 30 minutes before Lutathera infusion; continue during and for at least 3 hours after infusion. - Following Lutathera treatment, continue long-acting somatostatin analog every 4 weeks until disease progression or for 18 months following treatment initiation. <i>*Note: Lutathera is a radiopharmaceutical; handle with appropriate safety measures to minimize radiation exposure. Use waterproof gloves and effective radiation shielding when handling. Lutathera should be used by or under the control of healthcare providers who are qualified by specific training and experience in the safe use and handling of radiopharmaceuticals, and whose experience and training have been approved by the appropriate governmental agency authorized to license the use of radiopharmaceuticals.</i>

VI. Billing Code/Availability Information

HCPSC code:

- A9513 – Lutetium lu 177, dotatate, therapeutic, 1 millicurie; 1 billable unit = 1 millicurie

NDC:

- Lutathera 7.4 GBq (200 mCi) [370 MBq/mL (10 mCi/mL)] of lutetium Lu 177 dotatate single-dose vial: 69488-0003-XX

VII. References

- Lutathera [package insert]. Milburn, NJ; Advanced Accelerator Applications USA, Inc.; March 2023. Accessed June 2023.
- Strosberg J, El-Haddad G, Wolin E, et al. Phase 3 Trial of 177 Lu-Dotatate for Midgut Neuroendocrine Tumors. N Engl J Med. 2017 Jan 12;376(2):125-135.
- Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) lutetium lu 177 dotatate. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2023.
- Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for Neuroendocrine and Adrenal Tumors 2.2022. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN

LUTATHERA® (lutetium Lu 177 dotatate) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2023, Magellan Rx Management

Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc.” To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2023.

5. Brabander T, Van Der Zwan WA, Teunissen JJM, et al. Long-term efficacy, survival, and safety of [177Lu-DOTA0, Tyr3] octreotate in patients with gastroenteropancreatic and bronchial neuroendocrine tumors. Clin Cancer Res. 2017;23(16):4617-4624.
6. Kong G, Grozinsky-Glsberg S, Hofman MS, et al. Efficacy of peptide receptor radionuclide therapy for functional metastatic paraganglioma and pheochromocytoma. J Clin Endocrinol Metab 2017; 102(9): 3278-3287.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C74.10	Malignant neoplasm of medulla of unspecified adrenal gland
C74.11	Malignant neoplasm of medulla of right adrenal gland
C74.12	Malignant neoplasm of medulla of left adrenal gland
C74.90	Malignant neoplasm of unspecified part of unspecified adrenal gland
C74.91	Malignant neoplasm of unspecified part of right adrenal gland
C74.92	Malignant neoplasm of unspecified part of left adrenal gland
C75.5	Malignant neoplasm of aortic body and other paraganglia
C7A.00	Malignant carcinoid tumor of unspecified site
C7A.010	Malignant carcinoid tumor of the duodenum
C7A.011	Malignant carcinoid tumor of the jejunum
C7A.012	Malignant carcinoid tumor of the ileum
C7A.019	Malignant carcinoid tumor of the small intestine, unspecified portion
C7A.020	Malignant carcinoid tumor of the appendix
C7A.021	Malignant carcinoid tumor of the cecum
C7A.022	Malignant carcinoid tumor of the ascending colon
C7A.023	Malignant carcinoid tumor of the transverse colon
C7A.024	Malignant carcinoid tumor of the descending colon
C7A.025	Malignant carcinoid tumor of the sigmoid colon
C7A.026	Malignant carcinoid tumor of the rectum
C7A.029	Malignant carcinoid tumor of the large intestine, unspecified portion
C7A.090	Malignant carcinoid tumor of the bronchus and lung
C7A.091	Malignant carcinoid tumor of the thymus
C7A.092	Malignant carcinoid tumor of the stomach
C7A.093	Malignant carcinoid tumor of the kidney
C7A.094	Malignant carcinoid tumor of the foregut, unspecified
C7A.095	Malignant carcinoid tumor of the midgut, unspecified
C7A.096	Malignant carcinoid tumor of the hindgut, unspecified
C7A.098	Malignant carcinoid tumors of other sites

LUTATHERA® (lutetium Lu 177 dotatate) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2023, Magellan Rx Management

ICD-10	ICD-10 Description
C7A.8	Other malignant neuroendocrine tumors
C7B.00	Secondary carcinoid tumors, unspecified site
C7B.01	Secondary carcinoid tumors of distant lymph nodes
C7B.02	Secondary carcinoid tumors of liver
C7B.03	Secondary carcinoid tumors of bone
C7B.04	Secondary carcinoid tumors of peritoneum
C7B.09	Secondary carcinoid tumors of other sites
C7B.8	Other secondary neuroendocrine tumors
D3A.00	Benign carcinoid tumor of unspecified site
D3A.010	Benign carcinoid tumor of the duodenum
D3A.011	Benign carcinoid tumor of the jejunum
D3A.012	Benign carcinoid tumor of the ileum
D3A.019	Benign carcinoid tumor of the small intestine, unspecified portion
D3A.020	Benign carcinoid tumor of the appendix
D3A.021	Benign carcinoid tumor of the cecum
D3A.022	Benign carcinoid tumor of the ascending colon
D3A.023	Benign carcinoid tumor of the transverse colon
D3A.024	Benign carcinoid tumor of the descending colon
D3A.025	Benign carcinoid tumor of the sigmoid colon
D3A.026	Benign carcinoid tumor of the rectum
D3A.029	Benign carcinoid tumor of the large intestine, unspecified portion
D3A.090	Benign carcinoid tumor of the bronchus and lung
D3A.091	Benign carcinoid tumor of the thymus
D3A.092	Benign carcinoid tumor of the stomach
D3A.094	Benign carcinoid tumor of the foregut, unspecified
D3A.095	Benign carcinoid tumor of the midgut, unspecified
D3A.096	Benign carcinoid tumor of the hindgut, unspecified
D3A.098	Benign carcinoid tumors of other sites
E16.1	Other hypoglycemia
E16.3	Increased secretion of glucagon
E16.8	Other specified disorders of pancreatic internal secretion
E34.0	Carcinoid syndrome
Z85.020	Personal history of malignant carcinoid tumor of stomach
Z85.030	Personal history of malignant carcinoid tumor of large intestine
Z85.040	Personal history of malignant carcinoid tumor of rectum
Z85.060	Personal history of malignant carcinoid tumor of small intestine
Z85.07	Personal history of malignant neoplasm of pancreas
Z85.110	Personal history of malignant carcinoid tumor of bronchus and lung
Z85.230	Personal history of malignant carcinoid tumor of thymus

ICD-10	ICD-10 Description
Z85.858	Personal history of malignant neoplasm of other endocrine glands

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Articles (LCAs) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at:

<https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCA/LCD): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto Government Benefit Administrators, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC

Nondiscrimination & Language Access Policy

Aspirus Health Plan, Inc. complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, sex, sexual orientation, or gender identity. We do not exclude people or treat them differently because of race, color, national origin, age, disability, sex, sexual orientation, or gender identity.

We will:

Provide free aids and services to people with disabilities to communicate effectively with us, such as:

- Qualified sign language interpreters
- Written information in other formats (large print, audio, accessible electronic formats, other formats)

Provide free language services to people whose primary language is not English, such as:

- Qualified interpreters
- Information written in other languages

If you need these services, contact us at the phone number shown on the inside cover of this contract, your id card, or aspirushealthplan.com.

If you believe that we have failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, sex, sexual orientation, or gender identity, you can file a grievance with:

Nondiscrimination Grievance Coordinator
Aspirus Health Plan, Inc.
PO Box 1062
Minneapolis, MN 55440
Phone: 1.866.631.5404 (TTY: 711)
Fax: 763.847.4010
Email: customerservice@aspirushealthplan.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Nondiscrimination Grievance Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at:

U.S. Department of Health and Human Services
200 Independence Avenue, SW
Room 509F, HHH Building
Washington, D.C. 20201
1-800-368-1019, 800-537-7697 (TDD)

Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

Language Assistance Services

Albanian: KUJDES: Nëse flitni shqip, për ju ka në dispozicion shërbime të asistencës gjuhësore, pa pagesë. Telefononi në 1.866.631.5404 (TTY: 711).

Arabic: تنبيه: إذا كنت تتحدث اللغة العربية، فإن خدمات المساعدة اللغوية متاحة لك مجاناً. اتصل بن اعلی رقم الهاتف 1.866.631.5404 (رقم هاتف الصم والبك : 711)

French: ATTENTION : Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelezle 1.866.631.5404 (ATS : 711).

German: ACHTUNG: Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zur Verfügung. Rufnummer: 1.866.631.5404 (TTY: 711).

Hindi: _यान द_ : य_द आप िहंदी बोलते ह_ तो आपके िलए मु_त म_ भाषा सहायता सेवाएं उपल_ध ह_। 1.866.631.5404 (TTY: 711) पर कॉल कर_।

Hmong: LUS CEEV: Yog tias koj hais lus Hmoob, cov kev pab txog lus, muaj kev pab dawb rau koj. Hu rau 1.866.631.5404 (TTY: 711).

Korean: 주의: 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 1.866.631.5404 (TTY: 711)번으로 전화해 주십시오.

Polish: UWAGA: Jeżeli mówisz po polsku, możesz skorzystać z bezpłatnej pomocy językowej. Zadzwoń pod numer 1.866.631.5404 (TTY: 711).

Russian: ВНИМАНИЕ: Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода. Звоните 1.866.631.5404 (телетайп: 711).

Spanish: ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1.866.631.5404 (TTY: 711).

Tagalog: PAUNAWA: Kung nagsasalita ka ng Tagalog, maaari kang gumamit ng mga serbisyo ng tulong sa wika nangwalang bayad. Tumawag sa 1.866.631.5404 (TTY: 711)

Traditional Chinese: 注意: 如果您使用繁體中文, 您可以免費獲得語言援助服務。請致電 1.866.631.5404 (TTY: 711)。

Vietnamese: CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 1.866.631.5404 (TTY: 711).

Pennsylvania Dutch: Wann du Deitsch (Pennsylvania German / Dutch) schwetzscht, kannscht du mitaus Koschte ebbergricke, ass dihr helft mit die englisch Schprooch. Ruf selli Nummer uff: Call 1.866.631.5404 (TTY: 711).

Lao: ໄປດຊາຍ: ຖ້າວ່າ ທ່ານເວົ້າພາສາ ລາວ, ການບໍລິການຊ່ວຍເຫຼືອດ້ານພາສາ, ໂດຍບໍ່ເສັຽຄ່າ, ແມ່ນມີພ້ອມໃຫ້ທ່ານ. ໂທ 1.866.631.5404 (TTY: 711).