

# Simponi Aria (golimumab) injection, for intravenous use

Policy Number: MC/PC 039  
 Effective Date: December 1, 2025

[Instructions for Use](#)

Table of Contents	Page
<a href="#">Coverage Rationale</a> .....	1
<a href="#">Applicable Codes</a> .....	3
<a href="#">Background</a> .....	13
<a href="#">Clinical Evidence</a> .....	14
<a href="#">U.S. Food and Drug Administration</a> .....	16
<a href="#">References</a> .....	16
<a href="#">Policy History/Revision Information</a> .....	18
<a href="#">Instructions for Use</a> .....	18

**Related Policies**

- N/A

## Coverage Rationale

**This policy is applicable to Simponi Aria (golimumab) injection for intravenous infusion only.**

### Ankylosing Spondylitis

For initial coverage of Simponi Aria (golimumab) injection for adult patients with active Ankylosing Spondylitis (AS), the following will be required:

- Diagnosis of active ankylosing spondylitis **and**
- Minimum duration of one month trial and failure, contraindication, or intolerance to two different NSAIDs (e.g., ibuprofen, naproxen) at maximally tolerated doses **and**
- Prescribed by or in consultation with a rheumatologist

For reauthorization coverage of Simponi Aria (golimumab) injection for adult patients with active Ankylosing Spondylitis (AS), the following will be required:

- Patient demonstrates positive clinical response to therapy as evidenced by improvement from baseline for at least one of the following:
  - Disease activity (e.g., pain, fatigue, inflammation, stiffness)
  - Lab values (e.g., erythrocyte sedimentation rate, C-reactive protein level)
  - Function
  - Axial status (e.g., lumbar spine motion, chest expansion)
  - Total active joint count (swollen and tender)

### Polyarticular Juvenile Idiopathic Arthritis

For initial coverage of Simponi Aria (golimumab) injection for active polyarticular Juvenile Idiopathic Arthritis (pJIA) in patients 2 years of age and older, the following will be required:

- Diagnosis of moderately to severely active PJIA **and**

- Prescribed by or in consultation with a rheumatologist **and**
- Minimum duration of a 6-week trial and failure, contraindication, or in conventional therapies at maximally tolerated doses:
  - leflunomide
  - methotrexate

For reauthorization coverage of Simponi Aria (golimumab) injection for active polyarticular Juvenile Idiopathic Arthritis (pJIA) in patients 2 years of age and older, the following will be required:

- Patient demonstrates positive clinical response to therapy as evidenced by at least one of the following:
  - Reduction in the total active (swollen and tender) joint count from baseline
  - Improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline

### Psoriatic Arthritis

For initial coverage of Simponi Aria (golimumab) injection for active Psoriatic Arthritis (PsA) in patients 2 years of age and older, the following will be required:

- Diagnosis of active PsA **and**
- One of the following:
  - Actively inflamed joints
  - Dactylitis
  - Enthesitis
  - Axial disease
  - Active skin and/or nail involvement **and**
- Prescribed by or in consultation with one of the following:
  - Dermatologist
  - Rheumatologist

For reauthorization coverage of Simponi Aria (golimumab) injection for active Psoriatic Arthritis (PsA) in patients 2 years of age and older, the following will be required:

- Patient demonstrates positive clinical response to therapy as evidenced by at least one of the following:
  - Reduction in the total active (swollen and tender) joint count from baseline
  - Improvement in symptoms (e.g., pain, stiffness, pruritus, inflammation) from baseline

Reduction in body surface area (BSA) involvement from baseline

### Rheumatoid Arthritis (RA)

For initial coverage of Simponi Aria (golimumab) injection for adult patients with moderately to severely active Rheumatoid Arthritis (RA) in combination with methotrexate, the following will be required:

- Diagnosis of moderately to severely active RA **and**
- Minimum duration of a 3-month trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:
  - methotrexate
  - leflunomide
  - sulfasalazine **and**
- Used in combination with methotrexate **and**
- Prescribed by or in consultation with a rheumatologist

For reauthorization coverage of Simponi Aria (golimumab) injection for adult patients with moderately to severely active Rheumatoid Arthritis (RA) in combination with methotrexate, the following will be required:

- Patient demonstrates positive clinical response to therapy as evidenced by at least one of the following:
  - Reduction in the total active (swollen and tender) joint count from baseline **or**
  - Improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline

## Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

HCPCS Code	Description
J1602	Injection, golimumab, 1 mg, for intravenous use

ICD-10 Code	Description
L40.50	Arthropathic psoriasis, unspecified
L40.51	Distal interphalangeal psoriatic arthropathy
L40.52	Psoriatic arthritis mutilans
L40.53	Psoriatic spondylitis
L40.54	Psoriatic juvenile arthropathy
L40.59	Other psoriatic arthropathy
M05.00	Felty's syndrome, unspecified site
M05.011	Felty's syndrome, right shoulder
M05.012	Felty's syndrome, left shoulder
M05.019	Felty's syndrome, unspecified shoulder
M05.021	Felty's syndrome, right elbow
M05.022	Felty's syndrome, left elbow
M05.029	Felty's syndrome, unspecified elbow
M05.031	Felty's syndrome, right wrist
M05.032	Felty's syndrome, left wrist
M05.039	Felty's syndrome, unspecified wrist
M05.041	Felty's syndrome, right hand
M05.042	Felty's syndrome, left hand
M05.049	Felty's syndrome, unspecified hand
M05.051	Felty's syndrome, right hip
M05.052	Felty's syndrome, left hip
M05.059	Felty's syndrome, unspecified hip
M05.061	Felty's syndrome, right knee
M05.062	Felty's syndrome, left knee
M05.069	Felty's syndrome, unspecified knee
M05.071	Felty's syndrome, right ankle and foot
M05.072	Felty's syndrome, left ankle and foot
M05.079	Felty's syndrome, unspecified ankle and foot
M05.09	Felty's syndrome, multiple sites
M05.20	Rheumatoid vasculitis with rheumatoid arthritis of unspecified site

ICD-10 Code	Description
M05.211	Rheumatoid vasculitis with rheumatoid arthritis of right sh
M05.212	Rheumatoid vasculitis with rheumatoid arthritis of left shoulder
M05.219	Rheumatoid vasculitis with rheumatoid arthritis of unspecified shoulder
M05.221	Rheumatoid vasculitis with rheumatoid arthritis of right elbow
M05.222	Rheumatoid vasculitis with rheumatoid arthritis of left elbow
M05.229	Rheumatoid vasculitis with rheumatoid arthritis of unspecified elbow
M05.231	Rheumatoid vasculitis with rheumatoid arthritis of right wrist
M05.232	Rheumatoid vasculitis with rheumatoid arthritis of left wrist
M05.239	Rheumatoid vasculitis with rheumatoid arthritis of unspecified wrist
M05.241	Rheumatoid vasculitis with rheumatoid arthritis of right hand
M05.242	Rheumatoid vasculitis with rheumatoid arthritis of left hand
M05.249	Rheumatoid vasculitis with rheumatoid arthritis of unspecified hand
M05.251	Rheumatoid vasculitis with rheumatoid arthritis of right hip
M05.252	Rheumatoid vasculitis with rheumatoid arthritis of left hip
M05.259	Rheumatoid vasculitis with rheumatoid arthritis of unspecified hip
M05.261	Rheumatoid vasculitis with rheumatoid arthritis of right knee
M05.262	Rheumatoid vasculitis with rheumatoid arthritis of left knee
M05.269	Rheumatoid vasculitis with rheumatoid arthritis of unspecified knee
M05.271	Rheumatoid vasculitis with rheumatoid arthritis of right ankle and foot
M05.272	Rheumatoid vasculitis with rheumatoid arthritis of left ankle and foot
M05.279	Rheumatoid vasculitis with rheumatoid arthritis of unspecified ankle and foot
M05.29	Rheumatoid vasculitis with rheumatoid arthritis of multiple sites
M05.30	Rheumatoid heart disease with rheumatoid arthritis of unspecified site
M05.311	Rheumatoid heart disease with rheumatoid arthritis of right shoulder
M05.312	Rheumatoid heart disease with rheumatoid arthritis of left shoulder
M05.319	Rheumatoid heart disease with rheumatoid arthritis of unspecified shoulder
M05.321	Rheumatoid heart disease with rheumatoid arthritis of right elbow
M05.322	Rheumatoid heart disease with rheumatoid arthritis of left elbow
M05.329	Rheumatoid heart disease with rheumatoid arthritis of unspecified elbow
M05.331	Rheumatoid heart disease with rheumatoid arthritis of right wrist
M05.332	Rheumatoid heart disease with rheumatoid arthritis of left wrist
M05.339	Rheumatoid heart disease with rheumatoid arthritis of unspecified wrist
M05.341	Rheumatoid heart disease with rheumatoid arthritis of right hand
M05.342	Rheumatoid heart disease with rheumatoid arthritis of left hand
M05.349	Rheumatoid heart disease with rheumatoid arthritis of unspecified hand
M05.351	Rheumatoid heart disease with rheumatoid arthritis of right hip
M05.352	Rheumatoid heart disease with rheumatoid arthritis of left hip
M05.359	Rheumatoid heart disease with rheumatoid arthritis of unspecified hip
M05.361	Rheumatoid heart disease with rheumatoid arthritis of right knee
M05.362	Rheumatoid heart disease with rheumatoid arthritis of left knee

ICD-10 Code	Description
M05.369	Rheumatoid heart disease with rheumatoid arthritis of un
M05.371	Rheumatoid heart disease with rheumatoid arthritis of right ankle and foot
M05.372	Rheumatoid heart disease with rheumatoid arthritis of left ankle and foot
M05.379	Rheumatoid heart disease with rheumatoid arthritis of unspecified ankle and foot
M05.39	Rheumatoid heart disease with rheumatoid arthritis of multiple sites
M05.40	Rheumatoid myopathy with rheumatoid arthritis of unspecified site
M05.411	Rheumatoid myopathy with rheumatoid arthritis of right shoulder
M05.412	Rheumatoid myopathy with rheumatoid arthritis of left shoulder
M05.419	Rheumatoid myopathy with rheumatoid arthritis of unspecified shoulder
M05.421	Rheumatoid myopathy with rheumatoid arthritis of right elbow
M05.422	Rheumatoid myopathy with rheumatoid arthritis of left elbow
M05.429	Rheumatoid myopathy with rheumatoid arthritis of unspecified elbow
M05.431	Rheumatoid myopathy with rheumatoid arthritis of right wrist
M05.432	Rheumatoid myopathy with rheumatoid arthritis of left wrist
M05.439	Rheumatoid myopathy with rheumatoid arthritis of unspecified wrist
M05.441	Rheumatoid myopathy with rheumatoid arthritis of right hand
M05.442	Rheumatoid myopathy with rheumatoid arthritis of left hand
M05.449	Rheumatoid myopathy with rheumatoid arthritis of unspecified hand
M05.451	Rheumatoid myopathy with rheumatoid arthritis of right hip
M05.452	Rheumatoid myopathy with rheumatoid arthritis of left hip
M05.459	Rheumatoid myopathy with rheumatoid arthritis of unspecified hip
M05.461	Rheumatoid myopathy with rheumatoid arthritis of right knee
M05.462	Rheumatoid myopathy with rheumatoid arthritis of left knee
M05.469	Rheumatoid myopathy with rheumatoid arthritis of unspecified knee
M05.471	Rheumatoid myopathy with rheumatoid arthritis of right ankle and foot
M05.472	Rheumatoid myopathy with rheumatoid arthritis of left ankle and foot
M05.479	Rheumatoid myopathy with rheumatoid arthritis of unspecified ankle and foot
M05.49	Rheumatoid myopathy with rheumatoid arthritis of multiple sites
M05.50	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified site
M05.511	Rheumatoid polyneuropathy with rheumatoid arthritis of right shoulder
M05.512	Rheumatoid polyneuropathy with rheumatoid arthritis of left shoulder
M05.519	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified shoulder
M05.521	Rheumatoid polyneuropathy with rheumatoid arthritis of right elbow
M05.522	Rheumatoid polyneuropathy with rheumatoid arthritis of left elbow
M05.529	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified elbow
M05.531	Rheumatoid polyneuropathy with rheumatoid arthritis of right wrist
M05.532	Rheumatoid polyneuropathy with rheumatoid arthritis of left wrist
M05.539	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified wrist
M05.541	Rheumatoid polyneuropathy with rheumatoid arthritis of right hand
M05.542	Rheumatoid polyneuropathy with rheumatoid arthritis of left hand

ICD-10 Code	Description
M05.549	Rheumatoid polyneuropathy with rheumatoid arthritis of
M05.551	Rheumatoid polyneuropathy with rheumatoid arthritis of right hip
M05.552	Rheumatoid polyneuropathy with rheumatoid arthritis of left hip
M05.559	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified hip
M05.561	Rheumatoid polyneuropathy with rheumatoid arthritis of right knee
M05.562	Rheumatoid polyneuropathy with rheumatoid arthritis of left knee
M05.569	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified knee
M05.571	Rheumatoid polyneuropathy with rheumatoid arthritis of right ankle and foot
M05.572	Rheumatoid polyneuropathy with rheumatoid arthritis of left ankle and foot
M05.579	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified ankle and foot
M05.59	Rheumatoid polyneuropathy with rheumatoid arthritis of multiple sites
M05.60	Rheumatoid arthritis of unspecified site with involvement of other organs and systems
M05.611	Rheumatoid arthritis of right shoulder with involvement of other organs and systems
M05.612	Rheumatoid arthritis of left shoulder with involvement of other organs and systems
M05.619	Rheumatoid arthritis of unspecified shoulder with involvement of other organs and systems
M05.621	Rheumatoid arthritis of right elbow with involvement of other organs and systems
M05.622	Rheumatoid arthritis of left elbow with involvement of other organs and systems
M05.629	Rheumatoid arthritis of unspecified elbow with involvement of other organs and systems
M05.631	Rheumatoid arthritis of right wrist with involvement of other organs and systems
M05.632	Rheumatoid arthritis of left wrist with involvement of other organs and systems
M05.639	Rheumatoid arthritis of unspecified wrist with involvement of other organs and systems
M05.641	Rheumatoid arthritis of right hand with involvement of other organs and systems
M05.642	Rheumatoid arthritis of left hand with involvement of other organs and systems
M05.649	Rheumatoid arthritis of unspecified hand with involvement of other organs and systems
M05.651	Rheumatoid arthritis of right hip with involvement of other organs and systems
M05.652	Rheumatoid arthritis of left hip with involvement of other organs and systems
M05.659	Rheumatoid arthritis of unspecified hip with involvement of other organs and systems
M05.661	Rheumatoid arthritis of right knee with involvement of other organs and systems
M05.662	Rheumatoid arthritis of left knee with involvement of other organs and systems
M05.669	Rheumatoid arthritis of unspecified knee with involvement of other organs and systems
M05.671	Rheumatoid arthritis of right ankle and foot with involvement of other organs and systems
M05.672	Rheumatoid arthritis of left ankle and foot with involvement of other organs and systems
M05.679	Rheumatoid arthritis of unspecified ankle and foot with involvement of other organs and systems
M05.69	Rheumatoid arthritis of multiple sites with involvement of other organs and systems
M05.70	Rheumatoid arthritis with rheumatoid factor of unspecified site without organ or systems involvement
M05.711	Rheumatoid arthritis with rheumatoid factor of right shoulder without organ or systems involvement
M05.712	Rheumatoid arthritis with rheumatoid factor of left shoulder without organ or systems involvement
M05.719	Rheumatoid arthritis with rheumatoid factor of unspecified shoulder without organ or systems involvement

ICD-10 Code	Description
M05.721	Rheumatoid arthritis with rheumatoid factor of right elbow
M05.722	Rheumatoid arthritis with rheumatoid factor of left elbow without organ or systems involvement
M05.729	Rheumatoid arthritis with rheumatoid factor of unspecified elbow without organ or systems involvement
M05.731	Rheumatoid arthritis with rheumatoid factor of right wrist without organ or systems involvement
M05.732	Rheumatoid arthritis with rheumatoid factor of left wrist without organ or systems involvement
M05.739	Rheumatoid arthritis with rheumatoid factor of unspecified wrist without organ or systems involvement
M05.741	Rheumatoid arthritis with rheumatoid factor of right hand without organ or systems involvement
M05.742	Rheumatoid arthritis with rheumatoid factor of left hand without organ or systems involvement
M05.749	Rheumatoid arthritis with rheumatoid factor of unspecified hand without organ or systems involvement
M05.751	Rheumatoid arthritis with rheumatoid factor of right hip without organ or systems involvement
M05.752	Rheumatoid arthritis with rheumatoid factor of left hip without organ or systems involvement
M05.759	Rheumatoid arthritis with rheumatoid factor of unspecified hip without organ or systems involvement
M05.761	Rheumatoid arthritis with rheumatoid factor of right knee without organ or systems involvement
M05.762	Rheumatoid arthritis with rheumatoid factor of left knee without organ or systems involvement
M05.769	Rheumatoid arthritis with rheumatoid factor of unspecified knee without organ or systems involvement
M05.771	Rheumatoid arthritis with rheumatoid factor of right ankle and foot without organ or systems involvement
M05.772	Rheumatoid arthritis with rheumatoid factor of left ankle and foot without organ or systems involvement
M05.779	Rheumatoid arthritis with rheumatoid factor of unspecified ankle and foot without organ or systems involvement
M05.79	Rheumatoid arthritis with rheumatoid factor of multiple sites without organ or systems involvement
M05.7A	Rheumatoid arthritis with rheumatoid factor of other specified site without organ or systems involvement
M05.80	Other rheumatoid arthritis with rheumatoid factor of unspecified site
M05.811	Other rheumatoid arthritis with rheumatoid factor of right shoulder
M05.812	Other rheumatoid arthritis with rheumatoid factor of left shoulder
M05.819	Other rheumatoid arthritis with rheumatoid factor of unspecified shoulder
M05.821	Other rheumatoid arthritis with rheumatoid factor of right elbow
M05.822	Other rheumatoid arthritis with rheumatoid factor of left elbow
M05.829	Other rheumatoid arthritis with rheumatoid factor of unspecified elbow
M05.831	Other rheumatoid arthritis with rheumatoid factor of right wrist
M05.832	Other rheumatoid arthritis with rheumatoid factor of left wrist
M05.839	Other rheumatoid arthritis with rheumatoid factor of unspecified wrist
M05.841	Other rheumatoid arthritis with rheumatoid factor of right hand
M05.842	Other rheumatoid arthritis with rheumatoid factor of left hand

ICD-10 Code	Description
M05.849	Other rheumatoid arthritis with rheumatoid factor of unspecified site
M05.851	Other rheumatoid arthritis with rheumatoid factor of right hip
M05.852	Other rheumatoid arthritis with rheumatoid factor of left hip
M05.859	Other rheumatoid arthritis with rheumatoid factor of unspecified hip
M05.861	Other rheumatoid arthritis with rheumatoid factor of right knee
M05.862	Other rheumatoid arthritis with rheumatoid factor of left knee
M05.869	Other rheumatoid arthritis with rheumatoid factor of unspecified knee
M05.871	Other rheumatoid arthritis with rheumatoid factor of right ankle and foot
M05.872	Other rheumatoid arthritis with rheumatoid factor of left ankle and foot
M05.879	Other rheumatoid arthritis with rheumatoid factor of unspecified ankle and foot
M05.89	Other rheumatoid arthritis with rheumatoid factor of multiple sites
M05.8A	Other rheumatoid arthritis with rheumatoid factor of other specified site
M05.9	Rheumatoid arthritis with rheumatoid factor, unspecified
M06.00	Rheumatoid arthritis without rheumatoid factor, unspecified site
M06.011	Rheumatoid arthritis without rheumatoid factor, right shoulder
M06.012	Rheumatoid arthritis without rheumatoid factor, left shoulder
M06.019	Rheumatoid arthritis without rheumatoid factor, unspecified shoulder
M06.021	Rheumatoid arthritis without rheumatoid factor, right elbow
M06.022	Rheumatoid arthritis without rheumatoid factor, left elbow
M06.029	Rheumatoid arthritis without rheumatoid factor, unspecified elbow
M06.031	Rheumatoid arthritis without rheumatoid factor, right wrist
M06.032	Rheumatoid arthritis without rheumatoid factor, left wrist
M06.039	Rheumatoid arthritis without rheumatoid factor, unspecified wrist
M06.041	Rheumatoid arthritis without rheumatoid factor, right hand
M06.042	Rheumatoid arthritis without rheumatoid factor, left hand
M06.049	Rheumatoid arthritis without rheumatoid factor, unspecified hand
M06.051	Rheumatoid arthritis without rheumatoid factor, right hip
M06.052	Rheumatoid arthritis without rheumatoid factor, left hip
M06.059	Rheumatoid arthritis without rheumatoid factor, unspecified hip
M06.061	Rheumatoid arthritis without rheumatoid factor, right knee
M06.062	Rheumatoid arthritis without rheumatoid factor, left knee
M06.069	Rheumatoid arthritis without rheumatoid factor, unspecified knee
M06.071	Rheumatoid arthritis without rheumatoid factor, right ankle and foot
M06.072	Rheumatoid arthritis without rheumatoid factor, left ankle and foot
M06.079	Rheumatoid arthritis without rheumatoid factor, unspecified ankle and foot
M06.08	Rheumatoid arthritis without rheumatoid factor, vertebrae
M06.09	Rheumatoid arthritis without rheumatoid factor, multiple sites
M06.0A	Rheumatoid arthritis without rheumatoid factor, other specified site
M06.1	Adult-onset Still's disease
M06.20	Rheumatoid bursitis, unspecified site

ICD-10 Code	Description
M06.211	Rheumatoid bursitis, right shoulder
M06.212	Rheumatoid bursitis, left shoulder
M06.219	Rheumatoid bursitis, unspecified shoulder
M06.221	Rheumatoid bursitis, right elbow
M06.222	Rheumatoid bursitis, left elbow
M06.229	Rheumatoid bursitis, unspecified elbow
M06.231	Rheumatoid bursitis, right wrist
M06.232	Rheumatoid bursitis, left wrist
M06.239	Rheumatoid bursitis, unspecified wrist
M06.241	Rheumatoid bursitis, right hand
M06.242	Rheumatoid bursitis, left hand
M06.249	Rheumatoid bursitis, unspecified hand
M06.251	Rheumatoid bursitis, right hip
M06.252	Rheumatoid bursitis, left hip
M06.259	Rheumatoid bursitis, unspecified hip
M06.261	Rheumatoid bursitis, right knee
M06.262	Rheumatoid bursitis, left knee
M06.269	Rheumatoid bursitis, unspecified knee
M06.271	Rheumatoid bursitis, right ankle and foot
M06.272	Rheumatoid bursitis, left ankle and foot
M06.279	Rheumatoid bursitis, unspecified ankle and foot
M06.28	Rheumatoid bursitis, vertebrae
M06.29	Rheumatoid bursitis, multiple sites
M06.30	Rheumatoid nodule, unspecified site
M06.311	Rheumatoid nodule, right shoulder
M06.312	Rheumatoid nodule, left shoulder
M06.319	Rheumatoid nodule, unspecified shoulder
M06.321	Rheumatoid nodule, right elbow
M06.322	Rheumatoid nodule, left elbow
M06.329	Rheumatoid nodule, unspecified elbow
M06.331	Rheumatoid nodule, right wrist
M06.332	Rheumatoid nodule, left wrist
M06.339	Rheumatoid nodule, unspecified wrist
M06.341	Rheumatoid nodule, right hand
M06.342	Rheumatoid nodule, left hand
M06.349	Rheumatoid nodule, unspecified hand
M06.351	Rheumatoid nodule, right hip
M06.352	Rheumatoid nodule, left hip
M06.359	Rheumatoid nodule, unspecified hip
M06.361	Rheumatoid nodule, right knee

ICD-10 Code	Description
M06.362	Rheumatoid nodule, left knee
M06.369	Rheumatoid nodule, unspecified knee
M06.371	Rheumatoid nodule, right ankle and foot
M06.372	Rheumatoid nodule, left ankle and foot
M06.379	Rheumatoid nodule, unspecified ankle and foot
M06.38	Rheumatoid nodule, vertebrae
M06.39	Rheumatoid nodule, multiple sites
M06.80	Other specified rheumatoid arthritis, unspecified site
M06.811	Other specified rheumatoid arthritis, right shoulder
M06.812	Other specified rheumatoid arthritis, left shoulder
M06.819	Other specified rheumatoid arthritis, unspecified shoulder
M06.821	Other specified rheumatoid arthritis, right elbow
M06.822	Other specified rheumatoid arthritis, left elbow
M06.829	Other specified rheumatoid arthritis, unspecified elbow
M06.831	Other specified rheumatoid arthritis, right wrist
M06.832	Other specified rheumatoid arthritis, left wrist
M06.839	Other specified rheumatoid arthritis, unspecified wrist
M06.841	Other specified rheumatoid arthritis, right hand
M06.842	Other specified rheumatoid arthritis, left hand
M06.849	Other specified rheumatoid arthritis, unspecified hand
M06.851	Other specified rheumatoid arthritis, right hip
M06.852	Other specified rheumatoid arthritis, left hip
M06.859	Other specified rheumatoid arthritis, unspecified hip
M06.861	Other specified rheumatoid arthritis, right knee
M06.862	Other specified rheumatoid arthritis, left knee
M06.869	Other specified rheumatoid arthritis, unspecified knee
M06.871	Other specified rheumatoid arthritis, right ankle and foot
M06.872	Other specified rheumatoid arthritis, left ankle and foot
M06.879	Other specified rheumatoid arthritis, unspecified ankle and foot
M06.88	Other specified rheumatoid arthritis, vertebrae
M06.89	Other specified rheumatoid arthritis, multiple sites
M06.8A	Other specified rheumatoid arthritis, other specified site
M06.9	Rheumatoid arthritis, unspecified
M08.00	Unspecified juvenile rheumatoid arthritis of unspecified site
M08.011	Unspecified juvenile rheumatoid arthritis, right shoulder
M08.012	Unspecified juvenile rheumatoid arthritis, left shoulder
M08.019	Unspecified juvenile rheumatoid arthritis, unspecified shoulder
M08.021	Unspecified juvenile rheumatoid arthritis, right elbow
M08.022	Unspecified juvenile rheumatoid arthritis, left elbow
M08.029	Unspecified juvenile rheumatoid arthritis, unspecified elbow

ICD-10 Code	Description
M08.031	Unspecified juvenile rheumatoid arthritis, right wrist
M08.032	Unspecified juvenile rheumatoid arthritis, left wrist
M08.039	Unspecified juvenile rheumatoid arthritis, unspecified wrist
M08.041	Unspecified juvenile rheumatoid arthritis, right hand
M08.042	Unspecified juvenile rheumatoid arthritis, left hand
M08.049	Unspecified juvenile rheumatoid arthritis, unspecified hand
M08.051	Unspecified juvenile rheumatoid arthritis, right hip
M08.052	Unspecified juvenile rheumatoid arthritis, left hip
M08.059	Unspecified juvenile rheumatoid arthritis, unspecified hip
M08.061	Unspecified juvenile rheumatoid arthritis, right knee
M08.062	Unspecified juvenile rheumatoid arthritis, left knee
M08.069	Unspecified juvenile rheumatoid arthritis, unspecified knee
M08.071	Unspecified juvenile rheumatoid arthritis, right ankle and foot
M08.072	Unspecified juvenile rheumatoid arthritis, left ankle and foot
M08.079	Unspecified juvenile rheumatoid arthritis, unspecified ankle and foot
M08.08	Unspecified juvenile rheumatoid arthritis, vertebrae
M08.09	Unspecified juvenile rheumatoid arthritis, multiple sites
M08.0A	Unspecified juvenile rheumatoid arthritis, other specified site
M08.1	Juvenile ankylosing spondylitis
M08.20	Juvenile rheumatoid arthritis with systemic onset, unspecified site
M08.211	Juvenile rheumatoid arthritis with systemic onset, right shoulder
M08.212	Juvenile rheumatoid arthritis with systemic onset, left shoulder
M08.219	Juvenile rheumatoid arthritis with systemic onset, unspecified shoulder
M08.221	Juvenile rheumatoid arthritis with systemic onset, right elbow
M08.222	Juvenile rheumatoid arthritis with systemic onset, left elbow
M08.229	Juvenile rheumatoid arthritis with systemic onset, unspecified elbow
M08.231	Juvenile rheumatoid arthritis with systemic onset, right wrist
M08.232	Juvenile rheumatoid arthritis with systemic onset, left wrist
M08.239	Juvenile rheumatoid arthritis with systemic onset, unspecified wrist
M08.241	Juvenile rheumatoid arthritis with systemic onset, right hand
M08.242	Juvenile rheumatoid arthritis with systemic onset, left hand
M08.249	Juvenile rheumatoid arthritis with systemic onset, unspecified hand
M08.251	Juvenile rheumatoid arthritis with systemic onset, right hip
M08.252	Juvenile rheumatoid arthritis with systemic onset, left hip
M08.259	Juvenile rheumatoid arthritis with systemic onset, unspecified hip
M08.261	Juvenile rheumatoid arthritis with systemic onset, right knee
M08.262	Juvenile rheumatoid arthritis with systemic onset, left knee
M08.269	Juvenile rheumatoid arthritis with systemic onset, unspecified knee
M08.271	Juvenile rheumatoid arthritis with systemic onset, right ankle and foot
M08.272	Juvenile rheumatoid arthritis with systemic onset, left ankle and foot

ICD-10 Code	Description
M08.279	Juvenile rheumatoid arthritis with systemic onset, unspec
M08.28	Juvenile rheumatoid arthritis with systemic onset, vertebrae
M08.29	Juvenile rheumatoid arthritis with systemic onset, multiple sites
M08.2A	Juvenile rheumatoid arthritis with systemic onset, other specified site
M08.3	Juvenile rheumatoid polyarthritis (seronegative)
M08.80	Other juvenile arthritis, unspecified site
M08.811	Other juvenile arthritis, right shoulder
M08.812	Other juvenile arthritis, left shoulder
M08.819	Other juvenile arthritis, unspecified shoulder
M08.821	Other juvenile arthritis, right elbow
M08.822	Other juvenile arthritis, left elbow
M08.829	Other juvenile arthritis, unspecified elbow
M08.831	Other juvenile arthritis, right wrist
M08.832	Other juvenile arthritis, left wrist
M08.839	Other juvenile arthritis, unspecified wrist
M08.841	Other juvenile arthritis, right hand
M08.842	Other juvenile arthritis, left hand
M08.849	Other juvenile arthritis, unspecified hand
M08.851	Other juvenile arthritis, right hip
M08.852	Other juvenile arthritis, left hip
M08.859	Other juvenile arthritis, unspecified hip
M08.861	Other juvenile arthritis, right knee
M08.862	Other juvenile arthritis, left knee
M08.869	Other juvenile arthritis, unspecified knee
M08.871	Other juvenile arthritis, right ankle and foot
M08.872	Other juvenile arthritis, left ankle and foot
M08.879	Other juvenile arthritis, unspecified ankle and foot
M08.88	Other juvenile arthritis, vertebrae
M08.89	Other juvenile arthritis, multiple sites
M08.90	Juvenile arthritis, unspecified, unspecified site
M08.911	Juvenile arthritis, unspecified, right shoulder
M08.912	Juvenile arthritis, unspecified, left shoulder
M08.919	Juvenile arthritis, unspecified, unspecified shoulder
M08.921	Juvenile arthritis, unspecified, right elbow
M08.922	Juvenile arthritis, unspecified, left elbow
M08.929	Juvenile arthritis, unspecified, unspecified elbow
M08.931	Juvenile arthritis, unspecified, right wrist
M08.932	Juvenile arthritis, unspecified, left wrist
M08.939	Juvenile arthritis, unspecified, unspecified wrist
M08.941	Juvenile arthritis, unspecified, right hand

ICD-10 Code	Description
M08.942	Juvenile arthritis, unspecified, left hand
M08.949	Juvenile arthritis, unspecified, unspecified hand
M08.951	Juvenile arthritis, unspecified, right hip
M08.952	Juvenile arthritis, unspecified, left hip
M08.959	Juvenile arthritis, unspecified, unspecified hip
M08.961	Juvenile arthritis, unspecified, right knee
M08.962	Juvenile arthritis, unspecified, left knee
M08.969	Juvenile arthritis, unspecified, unspecified knee
M08.971	Juvenile arthritis, unspecified, right ankle and foot
M08.972	Juvenile arthritis, unspecified, left ankle and foot
M08.979	Juvenile arthritis, unspecified, unspecified ankle and foot
M08.98	Juvenile arthritis, unspecified, vertebrae
M08.99	Juvenile arthritis, unspecified, multiple sites
M08.9A	Juvenile arthritis, unspecified, other specified site
M45.0	Ankylosing spondylitis of multiple sites in spine
M45.1	Ankylosing spondylitis of occipito-atlanto-axial region
M45.2	Ankylosing spondylitis of cervical region
M45.3	Ankylosing spondylitis of cervicothoracic region
M45.4	Ankylosing spondylitis of thoracic region
M45.5	Ankylosing spondylitis of thoracolumbar region
M45.6	Ankylosing spondylitis lumbar region
M45.7	Ankylosing spondylitis of lumbosacral region
M45.8	Ankylosing spondylitis sacral and sacrococcygeal region
M45.9	Ankylosing spondylitis of unspecified sites in spine

## Background

Golimumab is a human IgG1  $\kappa$  monoclonal antibody specific for human tumor necrosis factor alpha (TNF  $\alpha$ ). TNF  $\alpha$  is an important mediator of the articular inflammation that is characteristic of these diseases and golimumab modulated the in vitro biological effects mediated by TNF in several bioassays, including the expression of adhesion proteins responsible for leukocyte infiltration (E-selectin, ICAM-1 and VCAM-1) and the secretion of proinflammatory cytokines (IL-6, IL-8, G-CSF and GM-CSF). Elevated TNF  $\alpha$  levels in the blood, synovium, and joints have been implicated in the pathophysiology of several chronic inflammatory diseases such as rheumatoid arthritis, psoriatic arthritis, and ankylosing spondylitis.

Golimumab was created using genetically engineered mice immunized with human TNF, resulting in an antibody with human-derived antibody variable and constant regions. It is produced by a recombinant cell line. Simponi Aria injection is a sterile solution of the golimumab antibody supplied in a 4-mL glass vial for intravenous infusion and was first approved by the FDA in April 2009 (Simponi Aria Prescribing Information 2025). The most common adverse reactions associated with Simponi Aria are upper respiratory tract infection, alanine aminotransferase increased, viral infection, aspartate aminotransferase increased, neutrophil count decreased, bronchitis, hypertension, and rash.

### **Ankylosing Spondylitis (AS)**

The FDA approval of Simponi (golimumab) for AS was based on a multicenter, randomized, double-blind, placebo-controlled trial in adult patients with active disease for at least 3 months (n = 356). Golimumab with or without a DMARD was compared to placebo with or without a DMARD and was found to significantly improve the signs and symptoms of AS demonstrated by the percentage of patients achieving an ASAS 20 response at week 14 (Inman et al 2008). Sustained improvements in ASAS 20 and ASAS 40 response rates were observed for up to 5 years in an open-label extension trial (Deodhar et al 2015). Safety profile through 5 years was consistent with other TNF inhibitors.

### **Polyarticular Juvenile Idiopathic Arthritis (pJIA)**

The approval of Simponi Aria (IV golimumab) for polyarticular JIA was based on an open-label Phase 3 study (n = 127). Children 2 to < 18 years of age with active polyarticular course JIA and inadequate response to MTX were enrolled. The primary endpoints were pharmacokinetic exposure and model-predicted steady-state area under the curve (AUCss) over an 8-week dosing interval at weeks 28 and 52. Other endpoints included ACR response rates. The ACR 30, 50, 70, and 90 response rates were 84%, 80%, 70%, and 47%, respectively, at week 28. Golimumab serum concentrations and AUCss were 0.40 mcg/mL and 399 mcg•day/mL at week 28. ACR response rates, serum concentrations, and AUCss were maintained at week 52 (Ruperto et al 2021).

### **Active Psoriatic Arthritis (PsA)**

The FDA approval of Simponi (golimumab) for PsA was based on the GO-REVEAL study, a multicenter, randomized, double-blind, placebo-controlled trial in adult patients with moderate to severely active PsA despite NSAID or DMARD therapy (n = 405). Golimumab with or without MTX compared to placebo with or without MTX, resulted in significant improvement in signs and symptoms as demonstrated by the percentage of patients achieving a ACR 20 response at week 14. The ACR responses observed in the golimumab-treated groups were similar in patients receiving and not receiving concomitant MTX therapy (Kavanaugh et al 2009).

Post-hoc analyses of the 5-year GO-REVEAL results evaluated the relationship between achieving minimal disease activity (MDA; defined as the presence of  $\geq 5$  of 7 PsA outcomes measures [ $\leq 1$  swollen joint,  $\leq 1$  tender joint, PASI  $\leq 1$ , patient pain score  $\leq 15$ , patient global disease activity score  $\leq 20$ , HAQ DI  $\leq 0.5$ , and  $\leq 1$  tender enthesis point]) and long-term radiographic outcomes including radiographic progression. Among golimumab-treated patients, achieving long-term MDA was associated with better long-term functional improvement, patient global assessment, and radiographic outcomes. Radiographic benefit was more pronounced in patients using MTX at baseline. The authors conclude that in patients with active PsA, aiming for MDA as part of a treat-to-target strategy may provide long-term functional and radiographic benefits (Kavanaugh et al 2016).

### **Rheumatoid Arthritis (RA)**

Simponi Aria (golimumab) was studied in 643 patients who received golimumab 2 mg/kg or 4 mg/kg intravenously (IV) every 12 weeks with or without MTX, or placebo with MTX. The proportion of patients meeting the primary endpoint of ACR 50 response was not significantly different between the golimumab with or without MTX groups and the placebo group. However, significantly more patients receiving golimumab plus MTX achieved an ACR 20 response at week 14 compared with patients receiving placebo plus MTX (53 vs 28%;  $p < 0.001$ ) (Kremer et al 2010). In the GO-FURTHER trial (n = 592), golimumab 2 mg/kg IV or placebo was given at weeks 0, 4 and then every 8 weeks. An increased percentage of patients treated with golimumab + MTX achieved ACR 20 response at week 14 (58.5% [231/395] of golimumab + MTX patients vs 24.9% [49/197] of placebo + MTX patients [ $p < 0.001$ ]) (Weinblatt et al 2013). In an open-label extension period, treatment was continued through week 100, with placebo-treated patients crossing over to golimumab at week 16 (early escape) or week 24. Clinical response was maintained through week 100, with an ACR 20 response of 68.1%. No unexpected AEs occurred (Bingham et al 2015).

## Clinical Guidelines

### **Ankylosing Spondylitis (AS)**

The American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network joint recommendations for treatment of AS and NRAS were updated in 2019. Patients with active AS or NRAS who do not respond to initial NSAID therapy are conditionally recommended to be treated with sulfasalazine, MTX, or tofacitinib; sulfasalazine or methotrexate should be considered only in patients with prominent peripheral arthritis or when TNF inhibitors are not available. Patients who do not respond to NSAID therapy are strongly recommended to receive treatment with a TNF inhibitor, although no particular TNF inhibitor is preferred. Treatment with a TNF inhibitor is conditionally recommended over tofacitinib, secukinumab, and ixekizumab in these patients. In patients with active disease who have primary nonresponse with a TNF inhibitor, treatment with secukinumab or ixekizumab is strongly recommended, and treatment with tofacitinib is conditionally recommended (Ward et al 2019).

Joint recommendations for the management of axial spondyloarthritis are available from ASAS and EULAR and were updated in 2022. The guideline notes that radiographic axial spondyloarthritis and non-radiographic axial spondyloarthritis are part of the same disease spectrum and therefore uses the term axial spondyloarthritis in recommendations. The guidelines state that NSAIDs should be used first-line in patients with pain and stiffness; other analgesics might be considered if NSAIDs have failed or are contraindicated or poorly tolerated. Glucocorticoid injections may be considered, but patients with axial disease should not receive long-term systemic glucocorticoids. Sulfasalazine may be considered in patients with peripheral arthritis, but patients with purely axial disease should normally not be treated with conventional DMARDs. TNF inhibitors, IL-17A inhibitors, or JAK inhibitors should be considered in patients with persistently high disease activity despite conventional treatments; current practice is to start with a TNF inhibitor or IL-17A inhibitor. In patients with a history of recurrent uveitis (UV) or active IBD, preference should be given to a monoclonal antibody against TNF. In patients with significant psoriasis, an IL-17 inhibitor may be preferred. Following failure of the first biologic or targeted synthetic DMARD, switching to another biologic DMARD (TNF inhibitor or IL-17A inhibitor) or a JAK inhibitor should be considered. For patients in sustained remission, tapering of a biologic DMARD can be considered (Ramiro et al 2023).

### **Polyarticular Juvenile Idiopathic Arthritis (pJIA)**

The ACR and Arthritis Foundation published a guideline for the treatment of JIA in 2019 focusing on therapy for non-systemic polyarthritis, sacroiliitis, and enthesitis. In children and adolescents with JIA and polyarthritis with moderate to high disease activity, addition of a biologic (TNF inhibitor, abatacept, or tocilizumab) is conditionally recommended. Patients with continued disease activity and primary TNF inhibitor failure are conditionally recommended to receive abatacept or tocilizumab over a second TNF inhibitor. Children and adolescents with JIA and active sacroiliitis despite treatment with NSAIDs are strongly recommended to add TNF inhibitor therapy over continuing NSAID monotherapy (Ringold et al 2019). A 2021 guideline from the ACR addresses the treatment of oligoarthritis, temporomandibular joint arthritis, and SJIA (Onel et al 2022).

### **Active Psoriatic Arthritis (PsA)**

EULAR 2019 PsA guidelines recommend biologic DMARDs in patients with peripheral arthritis and an inadequate response to at least 1 synthetic DMARD, such as MTX. For patients with peripheral arthritis, an inadequate response to at least 1 synthetic DMARD, and relevant skin involvement, biologics targeting IL-12/23 or IL-17 pathways may be considered.

The Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA) recommendations for PsA vary based on whether the arthritis is peripheral or axial and based on prior therapies, and may include DMARDs, NSAIDs, simple analgesics, a TNF inhibitor, inhibitors of IL-12/23 or IL-23, a JAK inhibitor, or a PDE-4 inhibitor (Coates et al 2022). The American College of Rheumatology/National Psoriasis Foundation guideline on PsA recommends that a TNF inhibitor is preferred in treatment-naïve patients with active PsA, although an oral therapy (MTX, sulfasalazine, leflunomide, cyclosporine, or apremilast) can be a first-line option in patients without severe PsA and without severe psoriasis, or if a patient has another compelling reason to avoid a TNF inhibitor. In patients who fail oral therapy, a

switch to a TNF inhibitor is preferred and placed ahead of IL-17 biologics (secukinumab, ixekicimab), abatacept, and tofacitinib (Singh et al 2019).

## Rheumatoid Arthritis (RA)

The American College of Rheumatology (ACR) recommends the use of conventional DMARDs, a TNF inhibitor, a non-TNF inhibitor biologic (tocilizumab, sarilumab, abatacept, or rituximab [only in patients that have had an inadequate response to TNF inhibitors or have a history of lymphoproliferative disorder]), or a JAK inhibitor (tofacitinib, baricitinib, upadacitinib). For patients who are not at target, switching to a medication in a different class is conditionally recommended over switching to a medication in the same class for patients receiving a biologic or JAK inhibitor. (Fraenkel et al 2021). EULAR guidelines for RA management were recently updated (Smolen et al 2023). EULAR recommends that therapy with DMARDs should be initiated as soon as the RA diagnosis is made with treatment aimed at reaching a target of sustained remission or low disease activity in every patient. If the treatment target is not achieved with the first conventional synthetic DMARD (csDMARD) strategy, in the absence of poor prognostic factors, other csDMARDs should be considered. If poor prognostic factors are present with csDMARD failure, a biological DMARD should be added; JAK inhibitors may be considered, but pertinent risk factors should be taken into account.

## U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

[Simponi Aria](#) is a tumor necrosis factor (TNF) blocker indicated for the treatment of:

- Adult patients with moderately to severely active Rheumatoid Arthritis (RA) in combination with methotrexate
- Active Psoriatic Arthritis (PsA) in patients 2 years of age and older
- Adult patients with active Ankylosing Spondylitis (AS)
- Active polyarticular Juvenile Idiopathic Arthritis (pJIA) in patients 2 years of age and older

## References

1. Bingham CO, Mendelsohn AM, Kim L, et al. Maintenance of clinical and radiographic benefit with intravenous golimumab therapy in patients with active rheumatoid arthritis despite methotrexate therapy: week-112 efficacy and safety results of the open-label long-term extension of a phase III, double-blind, randomized, placebo-controlled trial. *Arthritis Care Res*. 2015;67:1627-1636.
2. Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021. *Nat Rev Rheumatol*. 2022;18:465-479. doi:10.1038/s41584-022-00798-0
3. Deodhar A, Braun J, Inman RD, et al. Golimumab administered subcutaneously every 4 weeks in ankylosing spondylitis: 5-year results of the GO-RAISE study. *Ann Rheum Dis*. 2015;74:757-61.
4. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Rheumatol*. 2021;73(7):1108-1123. doi:10.1002/art.41752.
5. Inman R, Davis JC, Heijde DV et al. Efficacy and safety of golimumab in patients with ankylosing spondylitis: results of a randomized, double-blind, placebo-controlled, phase III trial. *Arthritis & Rheumatism: Official Journal of the American College of Rheumatology*. 2008;58(11):3402-12.
6. Kavanaugh A, Mease PJ, Gomez-Reino JJ, et al. Long-term (52-week) results of a phase III randomized, controlled trial of apremilast in patients with psoriatic arthritis. *J Rheumatol*. 2015;42:479-88.
7. Kavanaugh A, van der Heijde D, Beutler A, et al. Radiographic progression of patients with psoriatic arthritis who achieve minimal disease activity in response to golimumab therapy: results through 5 years of a randomized, placebo-controlled study. *Arthritis Care Res (Hoboken)*. 2016;68(2):267-74.

8. Kavanaugh A, van der Heijde D, Beutler A, et al. Radiographic progression and clinical outcomes in patients who achieve minimal disease activity in response to golimumab therapy: results of a placebo-controlled study. *Arthritis Care Res (Hoboken)*. 2016;68(2):267-74.
9. Kremer J, Ritchlin C, Mendelsohn A, et al. Golimumab, a new human anti-tumor necrosis factor alpha antibody, administered intravenously in patients with active rheumatoid arthritis. Forty-eight-week efficacy and safety results of a phase III randomized, double-blind, placebo-controlled study. *Arthritis Rheum*. 2010;62(4): 917–28.
10. Onel KB, Horton DB, Lovell DJ, et al. 2021 American College of Rheumatology guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for oligoarthritis, temporomandibular joint arthritis, and systemic juvenile idiopathic arthritis. *Arthritis Rheumatol*. 2022;74(4):553-569. doi:10.1002/art.42037.
11. Ramiro S, Nikiphorou E, Sepriano A, et al. ASAS-EULAR recommendations for the management of axial spondyloarthritis: 2022 update. *Ann Rheum Dis*. 2023;82:19-34. doi:10.1136/ard-2022-223296
12. Ringold S, Angeles-Han ST, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for non-systemic polyarthritis, sacroiliitis, and enthesitis. *Arthritis Care Res (Hoboken)*. 2019;71(6):717-734. doi: 10.1002/acr.23870.
13. Ruperto N, Brunner HI, Pacheco-Tena C, et al. Open-label phase 3 study of intravenous golimumab in patients with polyarticular juvenile idiopathic arthritis. *Rheumatology (Oxford)*. 2021;keab021. doi:10.1093/rheumatology/keab021.
14. Simponi Aria prescribing information. Janssen Biotech, Inc. Horsham, PA. September 2025. 3.
15. Singh JA, Guyatt G, Ogdie A, et al. Special Article: 2018 American College of Rheumatology/National Psoriasis Foundation guideline for the treatment of psoriatic arthritis. *Arthritis Rheumatol*. 2019;71(1):5-32. doi: 10.1002/art.40726.
16. Smolen JS, Beaulieu A, Rubbert-Roth A, et al. Effect of interleukin-6 receptor inhibition with tocilizumab in patients with rheumatoid arthritis (OPTION study): a double-blind, placebo-controlled, randomized trial. *Lancet*. 2008;371(9617):987-97.
17. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network recommendations for the treatment of ankylosing spondylitis and nonradiographic axial spondyloarthritis. *Arthritis Care Res (Hoboken)*. 2019;71(10):1285-1299. doi: 10.1002/acr.24025.
18. Weinblatt ME, Bingham CO, Mendelsohn AM, et al. Intravenous golimumab is effective in patients with active rheumatoid arthritis despite methotrexate therapy with responses as early as week 2: results of the phase 3, randomised, multicentre, double-blind, placebo-controlled GO-FURTHER trial. *Ann Rheum Dis*. 2013;72: 381-89.

## Policy History/Revision Information

Date	Summary of Changes
12/13/2023	Approved by OptumRx P&T Committee
11/21/2024	Annual Review. Updated references. Updated reauthorization criteria verbiage to "Patient demonstrates.."
11/20/2025	Annual review. Updated references and background section.

## Instructions for Use

This Medical Benefit Drug Policy provides assistance in interpreting standard benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard plan. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, please check the member specific benefit plan document and any applicable federal or state mandates. The insurance reserves the right to modify its Policies and Guidelines as necessary. This Medical Benefit Drug Policy is provided for informational purposes. It does not constitute medical advice.

OptumRx may also use tools developed by third parties to assist us in administering health benefits. OptumRx Medical Benefit Drug Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

# Nondiscrimination & Language Access Policy



Discrimination is Against the Law. Aspirus Health Plan, Inc. complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex, (including sex characteristics, including intersex traits; pregnancy or related conditions; sexual orientation, gender identity and sex stereotypes), consistent with the scope of sex discrimination described at 45 CFR § 92.101(a)(2). Aspirus Health Plan, Inc. does not exclude people or treat them less favorably because of race, color, national origin, age, disability, or sex.

Aspirus Health Plan, Inc.:

Provides people with disabilities reasonable modifications and free appropriate auxiliary aids and services to communicate effectively with us, such as:

- Qualified sign language interpreters.
- Written information in other formats (large print, audio, accessible electronic formats, other formats).

Provides free language assistance services to people whose primary language is not English, which may include:

- Qualified interpreters.
- Information written in other languages.

If you need reasonable modifications, appropriate auxiliary aids and services, or language assistance services, contact the Nondiscrimination Grievance Coordinator at the address, phone number, fax number, or email address below.

If you believe that Aspirus Health Plan, Inc. has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a *grievance* with:

Nondiscrimination Grievance Coordinator  
Aspirus Health Plan, Inc.  
PO Box 1890  
Southampton, PA 18966-9998  
Phone: 1-866-631-5404 (TTY: 711)  
Fax: 763-847-4010  
Email: customerservice@aspirushealthplan.com

You can file a *grievance* in person or by mail, fax, or email. If you need help filing a *grievance*, the Nondiscrimination Grievance Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at:

U.S. Department of Health and Human Services  
200 Independence Avenue, SW  
Room 509F, HHH Building  
Washington, D.C. 20201  
1.800.368.1019, 800.537.7697 (TDD)

Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>. This notice is available at Aspirus Health Plan, Inc.'s website: [https://aspirushealthplan.com/webdocs/70021-AHP-NonDiscrim\\_Lang-Assist-Notice.pdf](https://aspirushealthplan.com/webdocs/70021-AHP-NonDiscrim_Lang-Assist-Notice.pdf).

## Language Assistance Services

**Albanian:** KUJDES: Nëse flitmi shqip, për ju ka në dispozicion shërbime të asistencës gjuhësore, pa pagesë. Telefononi në 1-800-332-6501 (TTY: 711).

**Arabic:** تنبيه: إذا كنت تتحدث اللغة العربية، فإن خدمات المساعدة اللغوية متاحة لك مجاناً. اتصل بن اعلى رقم الهاتف 1-800-332-6501 (رقم هاتف الصم والبك : 711)

**French:** ATTENTION: Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelez le 1-800-332-6501 (ATS: 711).

**German:** ACHTUNG: Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zur Verfügung. Rufnummer: 1-800-332-6501 (TTY: 711).

**Hindi:** यान द : य द आप िहंदी बोलते ह तो आपके िलए मु त म भाषा सहायता सेवाएं उपल थ ह 1-800-332-6501 (TTY: 711) पर कॉल कर ।

**Hmong:** LUS CEEV: Yog tias koj hais lus Hmoob, cov kev pab txog lus, muaj kev pab dawb rau koj. Hu rau 1-800-332-6501 (TTY: 711).

**Korean:** 주의: 한국어를 사용하지는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 1-800-332-6501 (TTY: 711) 번으로 전화해 주십시오.

**Polish:** UWAGA: Jeżeli mówisz po polsku, możesz skorzystać z bezpłatnej pomocy językowej. Zadzwoń pod numer 1-800-332-6501 (TTY: 711).

**Russian:** ВНИМАНИЕ: Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода. Звоните 1-800-332-6501 (телетайп: 711).

**Spanish:** ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1-800-332-6501 (TTY: 711).

**Tagalog:** PAUNAWA: Kung nagsasalita ka ng Tagalog, maaari kang gumamit ng mga serbisyo ng tulong sa wika nangwalang bayad. Tumawag sa 1-800-332-6501 (TTY: 711).

**Traditional Chinese:** 注意：如果您使用繁體中文，您可以免費獲得語言援助服務。請致電 1-800-332-6501 (TTY: 711)

**Vietnamese:** CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 1-800-332-6501 (TTY: 711).

**Pennsylvania Dutch:** Wann du Deitsch (Pennsylvania German / Dutch) schwetzsch, kamscht du mitaus Koschte ebbergricke, ass dihr helft mit die englisch Schprooch. Ruf selli Nummer uff: Call 1-800-332-6501 (TTY: 711).

**Lao:** ໂປດຊາບ: ຖ້າວ່າ ທ່ານເວົ້າພາສາ ລາວ, ການບໍລິການຊ່ວຍເຫຼືອດ້ານພາສາ ໂດຍບໍ່ເສັຽຄ່າ, ຈະມີມື້ອມໃຫ້ທ່ານ. ໂທສ 1-800-332-6501 (TTY: 711).